

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 846 766 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
10.06.1998 Bulletin 1998/24

(21) Application number: 97307366.1

(22) Date of filing: 22.09.1997

(51) Int Cl.<sup>6</sup>: C12N 15/31, C07K 14/315,  
C07K 16/12, A61K 39/09,  
C12Q 1/68, G01N 33/50,  
G01N 33/68

(84) Designated Contracting States:  
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC  
NL PT SE

(30) Priority: 24.09.1996 US 27030 P  
10.03.1997 US 40656 P

(71) Applicants:  
• SMITHKLINE BEECHAM CORPORATION  
Philadelphia Pennsylvania 19103 (US)

• SMITHKLINE BEECHAM PLC  
Brentford, Middlesex TW8 9EP (GB)

(72) Inventor: Burnham, Martin K.R.,  
SmithKline Beecham Pharm.  
Collegeville, Pennsylvania 19426-0989 (US)

(74) Representative: Mallalieu, Catherine Louise et al  
D. Young & Co.,  
21 New Fetter Lane  
London EC4A 1DA (GB)

(54) **IgA Fc binding protein from Streptococcus pneumoniae**

(57) IgAFcBP polypeptides and DNA (RNA) encoding such IgAFcBP and a procedure for producing such polypeptides by recombinant techniques is disclosed. Also disclosed are methods for utilizing such IgAFcBP for the treatment of infection, and bacterial infections. Antagonists against such IgAFcBP and their use as a

therapeutic to treat infection and bacterial infections are also disclosed. Also disclosed are diagnostic assays for detecting diseases related to the presence of IgAFcBP nucleic acid sequences and the polypeptides in a host. Also disclosed are diagnostic assays for detecting polynucleotides encoding IgAFcBP and for detecting the polypeptide in a host.

EP 0 846 766 A2

## Description

This invention relates, in part, to newly identified polynucleotides and polypeptides; variants and derivatives of these polynucleotides and polypeptides; processes for making these polynucleotides and these polypeptides, and their variants and derivatives; agonists and antagonists of the polypeptides; and uses of these polynucleotides, polypeptides, variants, derivatives, agonists and antagonists. This invention also relates to inhibiting the biosynthesis or action of such polynucleotides or polypeptides and to the use of such inhibitors in therapy. In particular, in these and in other regards, the invention relates to polynucleotides and polypeptides of IgA Fc binding protein, hereinafter referred to as "IgAFcBP".

The Streptococci make up a medically important genera of microbes known to cause several types of disease in humans, including otitis media, pneumonia and meningitis. Since its isolation more than 100 years ago, *Streptococcus pneumoniae* has been one of the more intensively studied microbes. For example, much of our early understanding that DNA is, in fact, the genetic material was predicated on the work of Griffith and of Avery, Macleod and McCarty using this microbe. Despite the vast amount of research with *S. pneumoniae*, many questions concerning the virulence of this microbe remain.

While certain Streptococcal factors associated with pathogenicity have been identified, e.g., capsule polysaccharides, peptidoglycans, pneumolysins, PspA Complement factor H binding component, autolysin, neuraminidase, peptidase permeases, hydrogen peroxide, IgA1 protease, the list is certainly not complete. Further very little is known concerning the temporal expression of such genes during infection and disease progression in a mammalian host. Discerning the sets of genes the bacterium is likely to be expressing at the different stages of infection, particularly when an infection is established, provides critical information for the screening and characterization of novel antibacterials which can interrupt pathogenesis. In addition to providing a fuller understanding of known proteins, such an approach will identify previously unrecognised targets.

Several species of gram-positives pathogenic bacteria express cell wall proteins that bind to the Fc region of IgA in humans. Two of these from Streptococcus have been studied in detail: protein Arp from the group A Streptococcus and protein Bac from the group B Streptococcus (see Lindahl, G., Akerstrom, B., Stenberg, L., Frithz, E and Heden, L. O. (1991) in *Genetics and Molecular Biology of Streptococci, Lactococci and Enterococci* (Dunni, GM et al., eds.) pp155-159, American Soc. Microbiol. and also Jerlstrom, P.G., Chhatwal, G.S. and Timmis, K.N. (1991) *Mol. Microbiol.* 5: 843). Such proteins are valuable as immunochemical tools as it is likely that they are used by the bacteria to evade the defense mechanisms of the infected host (see Stenberg L., et al. (1994) *J. Biol. Chem.* 269(18): 13458).

This invention provides a IgA Fc region binding protein from *Streptococcus pneumoniae*, strain 0100993, comprising the polypeptide sequence given herein, or polypeptide fragments, analogues or derivatives thereof homologous at the amino acid level to the Streptococcus B Bac protein.

Clearly, there is a need for factors that may be used to screen compounds for antibacterial activity and which may also be used to determine their roles in pathogenesis of infection, dysfunction and disease. There is a need, therefore, for identification and characterization of such factors which can play a role in preventing, ameliorating or correcting infections, dysfunctions or diseases.

The polypeptide of the present invention has amino acid sequence homology to known streptococcal binding protein.

## SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the present invention to provide polypeptides, *inter alia*, that have been identified as novel IgAFcBP by homology between the amino acid sequence set out in Figure 2 and known amino acid sequences of other proteins such as streptococcal binding proteins.

It is a further object of the invention, moreover, to provide polynucleotides that encode IgAFcBP, particularly polynucleotides that encode the polypeptide herein designated IgAFcBP.

In a particularly preferred embodiment of this aspect of the invention the polynucleotide comprises the region encoding IgAFcBP in the sequence set out in Figure 1.

In another particularly preferred embodiment of the present invention there is a novel cell surface protein from *Streptococcus pneumoniae* comprising the amino acid sequence of SEQ ID NO:2, or a fragment, analogue or derivative thereof.

In accordance with this aspect of the present invention there is provided an isolated nucleic acid molecule encoding a mature polypeptide expressible by the *Streptococcus pneumoniae* polynucleotide contained in NCIMB Deposit No. 40794.

In accordance with this aspect of the invention there are provided isolated nucleic acid molecules encoding IgAFcBP, particularly *Streptococcus* IgAFcBP, including mRNAs, cDNAs, genomic DNAs and, in further embodiments of this aspect of the invention, biologically, diagnostically, clinically or therapeutically useful variants, analogs or deriv-

atives thereof, or fragments thereof, including fragments of the variants, analogs and derivatives.

Among the particularly preferred embodiments of this aspect of the invention are naturally occurring allelic variants of IgAFcBP.

5 In accordance with this aspect of the invention there are provided novel polypeptides of *Streptococcus* origin referred to herein as IgAFcBP as well as biologically, diagnostically or therapeutically useful fragments, variants and derivatives thereof, variants and derivatives of the fragments, and analogs of the foregoing.

10 It is also an object of the invention to provide IgAFcBP polypeptides, particularly IgAFcBP polypeptides, that may be employed for therapeutic purposes, for example, to treat disease, including treatment by conferring host immunity against infections, such as bacterial infections and particularly streptococcal infections, such as those of *Streptococcus pneumoniae*.

In accordance with yet a further aspect of the present invention, there is provided the use of a polypeptide of the invention for therapeutic or prophylactic purposes, for example, as an antibacterial agent or a vaccine.

In accordance with another aspect of the present invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization.

15 Among the particularly preferred embodiments of this aspect of the invention are variants of IgAFcBP polypeptide encoded by naturally occurring alleles of the IgAFcBP gene.

It is another object of the invention to provide a process for producing the aforementioned polypeptides, polypeptide fragments, variants and derivatives, fragments of the variants and derivatives, and analogs of the foregoing.

20 In a preferred embodiment of this aspect of the invention there are provided methods for producing the aforementioned IgAFcBP polypeptides comprising culturing host cells having expressibly incorporated therein an exogenously-derived IgAFcBP-encoding polynucleotide under conditions for expression of IgAFcBP in the host and then recovering the expressed polypeptide.

25 In accordance with another object the invention there are provided products, compositions, processes and methods that utilize the aforementioned polypeptides and polynucleotides, *inter alia* for research, biological, clinical and therapeutic purposes.

In accordance with yet another aspect of the present invention, there are provided inhibitors to such polypeptides, useful as antibacterial agents. In particular, there are provided antibodies against such polypeptides.

30 In accordance with certain preferred embodiments of this aspect of the invention, there are provided products, compositions and methods, *inter alia*: assessing IgAFcBP expression in cells by detecting IgAFcBP polypeptides or IgAFcBP-encoding mRNA; to treat bacterial infections, such as streptococcal infections, *in vitro*, *ex vivo* or *in vivo* by exposing cells to IgAFcBP polypeptides or polynucleotides as disclosed herein; assaying genetic variation and aberrations, such as defects, in IgAFcBP genes; and administering a IgAFcBP polypeptide or polynucleotide to an organism to raise an immunological response against a bacteria, especially a *Streptococcus*.

35 In accordance with certain preferred embodiments of this and other aspects of the invention there are probes that hybridize to IgAFcBP sequences.

In certain additional preferred embodiments of this aspect of the invention there are provided antibodies against IgAFcBP polypeptides. In certain particularly preferred embodiments in this regard, the antibodies are highly selective for IgAFcBP.

40 In accordance with another aspect of the present invention, there are provided IgAFcBP agonists. Among preferred agonists are molecules that mimic IgAFcBP, that bind to IgAFcBP-binding molecules, and that elicit or augment IgAFcBP-induced responses. Also among preferred agonists are molecules that interact with IgAFcBP or IgAFcBP polypeptides, or with other modulators of IgAFcBP activities, and thereby potentiate or augment an effect of IgAFcBP or more than one effect of IgAFcBP and are which are also preferably bacteriostatic or bacteriocidal.

45 In accordance with yet another aspect of the present invention, there are provided IgAFcBP antagonists. Among preferred antagonists are those which mimic IgAFcBP so as to bind to IgAFcBP-binding molecules but not elicit a IgAFcBP-induced response or more than one IgAFcBP-induced response. Also among preferred antagonists are molecules that bind to or interact with IgAFcBP so as to inhibit an effect of IgAFcBP or more than one effect of IgAFcBP or which prevent expression of IgAFcBP.

50 In a further aspect of the invention there are provided compositions comprising a IgAFcBP polynucleotide or a IgAFcBP polypeptide for administration to cells *in vitro*, to cells *ex vivo* and to cells *in vivo*, or to a multicellular organism. In certain preferred embodiments of this aspect of the invention, the compositions comprise a IgAFcBP polynucleotide for expression of a IgAFcBP polypeptide in a host organism to raise an immunological response, preferably to raise immunity in such host against *Streptococcus* or related organisms.

55 Other objects, features, advantages and aspects of the present invention will become apparent to those of skill from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

## BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings depict certain embodiments of the invention. They are illustrative only and do not limit the invention otherwise disclosed herein.

Figure 1 shows the polynucleotide sequence of *Streptococcus* IgAfCBP [SEQ ID NO:1].

Figure 2 shows the amino acid sequence of IgAfCBP [SEQ ID NO:2] deduced from the polynucleotide sequence of Figure 1.

## GLOSSARY

The following illustrative explanations are provided to facilitate understanding of certain terms used frequently herein, particularly in the Examples. The explanations are provided as a convenience and are not limitative of the invention.

IgAfCBP-BINDING MOLECULE, as used herein, refers to molecules or ions which bind or interact specifically with IgAfCBP polypeptides or polynucleotides of the present invention, including, for example enzyme substrates, cell membrane components and classical receptors. Binding between polypeptides of the invention and such molecules, including binding or binding or interaction molecules may be exclusive to polypeptides of the invention, which is preferred, or it may be highly specific for polypeptides of the invention, which is also preferred, or it may be highly specific to a group of proteins that includes polypeptides of the invention, which is preferred, or it may be specific to several groups of proteins at least one of which includes a polypeptide of the invention. Binding molecules also include antibodies and antibody-derived reagents that bind specifically to polypeptides of the invention.

DIGESTION of DNA refers to cleavage of DNA, for example, with a restriction enzyme that acts only at certain sequences in the DNA. The various restriction enzymes referred to herein are commercially available and their reaction conditions, cofactors and other requirements for use are well known and routine to the skilled artisan. For analytical purposes, typically, 1 µg of plasmid or DNA fragment is digested with about 2 units of enzyme in about 20 microliter of reaction buffer. For the purpose of isolating DNA fragments for plasmid construction, typically 5 to 50 microgram of DNA are digested with 20 to 250 units of enzyme in proportionately larger volumes. Appropriate buffers and substrate amounts for particular restriction enzymes are described in standard laboratory manuals, such as those referenced below, and they are specified by commercial suppliers. Incubation times of about 1 hour at 37°C are ordinarily used, but conditions may vary in accordance with standard procedures, the supplier's instructions and the particulars of the reaction. After digestion, reactions may be analyzed, and fragments may be purified by electrophoresis through an agarose or polyacrylamide gel, using well known methods that are routine for those skilled in the art.

GENETIC ELEMENT generally means a polynucleotide comprising a region that encodes a polypeptide or a polynucleotide region that regulates replication, transcription or translation or other processes important to expression of the polypeptide in a host cell, or a polynucleotide comprising both a region that encodes a polypeptide and a region operably linked thereto that regulates expression. Genetic elements may be comprised within a vector that replicates as an episomal element; that is, as a molecule physically independent of the host cell genome. They may be comprised within plasmids. Genetic elements also may be comprised within a host cell genome; not in their natural state but, rather, following manipulation such as isolation, cloning and introduction into a host cell in the form of purified DNA or in a vector, among others.

HOST CELL is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

IDENTITY or SIMILARITY, as known in the art, are relationships between two polypeptide sequences or two polynucleotide sequences, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between two polypeptide or two polynucleotide sequences as determined by the match between two strings of such sequences. Both identity and similarity can be readily calculated (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D. W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity and similarity between two polynucleotide or two polypeptide sequences, both terms are well known to skilled artisans (*Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to those disclosed in Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48:1073 (1988). Preferred methods to determine identity are designed to give the largest match between the two sequences tested. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include,

but are not limited to, GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Altschul, S.F. et al., *J. Molec. Biol.* 215: 403 (1990)).

ISOLATED means altered "by the hand of man" from its natural state; i.e., that, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a naturally occurring polynucleotide or a polypeptide naturally present in a living organism in its natural state is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. For example, with respect to polynucleotides, the term isolated means that it is separated from the chromosome and cell in which it naturally occurs. As part of or following isolation, such polynucleotides can be joined to other polynucleotides, such as DNAs, for mutagenesis, to form fusion proteins, and for propagation or expression in a host, for instance. The isolated polynucleotides, alone or joined to other polynucleotides such as vectors, can be introduced into host cells, in culture or in whole organisms. Introduced into host cells in culture or in whole organisms, such DNAs still would be isolated, as the term is used herein, because they would not be in their naturally occurring form or environment. Similarly, the polynucleotides and polypeptides may occur in a composition, such as a media formulations, solutions for introduction of polynucleotides or polypeptides, for example, into cells, compositions or solutions for chemical or enzymatic reactions, for instance, which are not naturally occurring compositions, and, therein remain isolated polynucleotides or polypeptides within the meaning of that term as it is employed herein.

LIGATION refers to the process of forming phosphodiester bonds between two or more polynucleotides, which most often are double stranded DNAs. Techniques for ligation are well known to the art and protocols for ligation are described in standard laboratory manuals and references, such as, for instance, Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989) and Maniatis et al., pg. 146, as cited below.

OLIGONUCLEOTIDE(S) refers to short polynucleotides. Often the term refers to single-stranded deoxyribonucleotides, but it can refer as well to single- or double-stranded ribonucleotides, RNA:DNA hybrids and double-stranded DNAs, among others. Oligonucleotides, such as single-stranded DNA probe oligonucleotides, often are synthesized by chemical methods, such as those implemented on automated oligonucleotide synthesizers. However, oligonucleotides can be made by a variety of other methods, including *in vitro* recombinant DNA-mediated techniques and by expression of DNAs in cells and organisms. Initially, chemically synthesized DNAs typically are obtained without a 5' phosphate. The 5' ends of such oligonucleotides are not substrates for phosphodiester bond formation by ligation reactions that employ DNA ligases typically used to form recombinant DNA molecules. Where ligation of such oligonucleotides is desired, a phosphate can be added by standard techniques, such as those that employ a kinase and ATP. The 3' end of a chemically synthesized oligonucleotide generally has a free hydroxyl group and, in the presence of a ligase, such as T4 DNA ligase, readily will form a phosphodiester bond with a 5' phosphate of another polynucleotide, such as another oligonucleotide. As is well known, this reaction can be prevented selectively, where desired, by removing the 5' phosphates of the other polynucleotide(s) prior to ligation.

PLASMIDS generally are designated herein by a lower case p preceded and/or followed by capital letters and/or numbers, in accordance with standard naming conventions that are familiar to those of skill in the art. Starting plasmids disclosed herein are either commercially available, publicly available, or can be constructed from available plasmids by routine application of well known, published procedures. Many plasmids and other cloning and expression vectors that can be used in accordance with the present invention are well known and readily available to those of skill in the art. Moreover, those of skill readily may construct any number of other plasmids suitable for use in the invention. The properties, construction and use of such plasmids, as well as other vectors, in the present invention will be readily apparent to those of skill from the present disclosure.

POLYNUCLEOTIDE(S) generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. Thus, for instance, polynucleotides as used herein refers to, among others, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded, or a mixture of single- and double-stranded regions. In addition, polynucleotide as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term polynucleotide includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotides" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term polynucleotide as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA

characteristic of viruses and cells, including simple and complex cells, *inter alia*.

POLYPEPTIDES, as used herein, includes all polypeptides as described below. The basic structure of polypeptides is well known and has been described in innumerable textbooks and other publications in the art. In this context, the term is used herein to refer to any peptide or protein comprising two or more amino acids joined to each other in a linear chain by peptide bonds. As used herein, the term refers to both short chains, which also commonly are referred to in the art as peptides, oligopeptides and oligomers, for example, and to longer chains, which generally are referred to in the art as proteins, of which there are many types. It will be appreciated that polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids, and that many amino acids, including the terminal amino acids, may be modified in a given polypeptide, either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques which are well known to the art. Even the common modifications that occur naturally in polypeptides are too numerous to list exhaustively here, but they are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. Among the known modifications which may be present in polypeptides of the present are, to name an illustrative few, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. Such modifications are well known to those of skill and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as, for instance *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as, for example, those provided by Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). It will be appreciated, as is well known and as noted above, that polypeptides are not always entirely linear. For instance, polypeptides may be branched as a result of ubiquitination, and they may be circular, with or without branching, generally as a result of posttranslation events, including natural processing event and events brought about by human manipulation which do not occur naturally. Circular, branched and branched circular polypeptides may be synthesized by non-translation natural process and by entirely synthetic methods, as well. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. In fact, blockage of the amino or carboxyl group in a polypeptide, or both, by a covalent modification, is common in naturally occurring and synthetic polypeptides and such modifications may be present in polypeptides of the present invention, as well. For instance, the amino terminal residue of polypeptides made in *E. coli* or other cells, prior to proteolytic processing, almost invariably will be N-formylmethionine. During post-translational modification of the peptide, a methionine residue at the NH<sub>2</sub>-terminus may be deleted. Accordingly, this invention contemplates the use of both the methionine-containing and the methionineless amino terminal variants of the protein of the invention. The modifications that occur in a polypeptide often will be a function of how it is made. For polypeptides made by expressing a cloned gene in a host, for instance, the nature and extent of the modifications in large part will be determined by the host cell posttranslational modification capacity and the modification signals present in the polypeptide amino acid sequence. For instance, as is well known, glycosylation often does not occur in bacterial hosts such as, for example, *E. coli*. Accordingly, when glycosylation is desired, a polypeptide should be expressed in a glycosylating host, generally a eukaryotic cell. Insect cell often carry out the same posttranslational glycosylations as mammalian cells and, for this reason, insect cell expression systems have been developed to express efficiently mammalian proteins having native patterns of glycosylation, *inter alia*. Similar considerations apply to other modifications. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. In general, as used herein, the term polypeptide encompasses all such modifications, particularly those that are present in polypeptides synthesized by expressing a polynucleotide in a host cell.

TRANSFORMATION is the process by which a cell is "transformed" by exogenous DNA when such exogenous DNA has been introduced inside the cell membrane. Exogenous DNA may or may not be integrated (covalently linked) into chromosomal DNA making up the genome of the cell. In prokaryotes and yeasts, for example, the exogenous DNA may be maintained on an episomal element, such as a plasmid. With respect to eukaryotic cells, a stably transformed or transfected cell is one in which the exogenous DNA has become integrated into the chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to

establish cell lines or clones comprised of a population of daughter cell containing the exogenous DNA.

VARIANT(S) of polynucleotides or polypeptides, as the term is used herein, are polynucleotides or polypeptides that differ from a reference polynucleotide or polypeptide, respectively. Variants in this sense are described below and elsewhere in the present disclosure in greater detail. (1) A polynucleotide that differs in nucleotide sequence from another, reference polynucleotide. Generally, differences are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical. As noted below, changes in the nucleotide sequence of the variant may be silent. That is, they may not alter the amino acids encoded by the polynucleotide. Where alterations are limited to silent changes of this type a variant will encode a polypeptide with the same amino acid sequence as the reference. Also as noted below, changes in the nucleotide sequence of the variant may alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Such nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. (2) A polypeptide that differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference and the variant are closely similar overall and, in many region, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions, fusions and truncations, which may be present in any combination.

## DESCRIPTION OF THE INVENTION

The present invention relates to novel IgAFcBP polypeptides and polynucleotides, among other things, as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of a novel IgAFcBP gene of *Streptococcus pneumoniae*, which is related by amino acid sequence homology to streptococcal IgAFcBP polypeptides. The invention relates especially to IgAFcBP having the nucleotide and amino acid sequences set out in Figure 1 and Figure 2 respectively, and to the IgAFcBP nucleotide and amino acid sequences of the DNA in NCIMB Deposit No. 40794, which is herein referred to as "the deposited clone" or as the "DNA of the deposited clone." It will be appreciated that the nucleotide and amino acid sequences set out in Figures 1 [SEQ ID NO: 1] and 2 [SEQ ID NO:2] were obtained by sequencing the DNA of the deposited clone. Hence, the sequence of the deposited clone is controlling as to any discrepancies between it (and the sequence it encodes) and the sequences of Figure 1 [SEQ ID NO: 1] and Figure 2 [SEQ ID NO:2].

Recently techniques have become available to evaluate temporal gene expression in bacteria, particularly as it applies to viability under laboratory and host infection conditions. A number of methods can be used to identify genes which are essential to survival *per se*, or essential to the establishment/maintenance of an infection. Identification of expression of a sequence by one of these methods yields additional information about its function and permits the selection of such sequence for further development as a screening target. Briefly, these approaches include:

### 1) Signature Tagged Mutagenesis (STM):

This technique is described by Hensel *et al.*, *Science* 269: 400-403(1995), the contents of which is incorporated by reference for background purposes. Signature tagged mutagenesis identifies genes necessary for the establishment/maintenance of infection in a given infection model.

The basis of the technique is the random mutagenesis of target organism by various means (e.g., transposons) such that unique DNA sequence tags are inserted in close proximity to the site of mutation. The tags from a mixed population of bacterial mutants and bacteria recovered from an infected hosts are detected by amplification, radiolabeling and hybridisation analysis. Mutants attenuated in virulence are revealed by absence of the tag from the pool of bacteria recovered from infected hosts.

In *Streptococcus pneumoniae*, because the transposon system is less well developed, a more efficient way of creating the tagged mutants is to use the insertion-duplication mutagenesis technique as described by Morrison *et al.*, *J. Bacteriol.* 159:870 (1984) the contents of which is incorporated by reference for background purposes.

### 2) In Vivo Expression Technology (IVET):

This technique is described by Camilli *et al.*, *Proc. Nat'l. Acad. Sci. USA* 91:2634-2638 (1994), the contents of which is incorporated by reference for background purposes. IVET identifies genes up-regulated during infection when compared to laboratory cultivation, implying an important role in infection. Sequences identified by this technique are implied to have a significant role in infection establishment/maintenance.

In this technique random chromosomal fragments of target organism are cloned upstream of a promoter-less recombinase gene in a plasmid vector. This construct is introduced into the target organism which carries an antibacterial resistance gene flanked by resolvase sites. Growth in the presence of the antibacterial removes from the population those fragments cloned into the plasmid vector capable of supporting transcription of the recombinase gene and



therefore have caused loss of antibacterial resistance. The resistant pool is introduced into a host and at various times after infection bacteria may be recovered and assessed for the presence of antibacterial resistance. The chromosomal fragment carried by each antibacterial sensitive bacterium should carry a promoter or portion of a gene normally up-regulated during infection. Sequencing upstream of the recombinase gene allows identification of the up regulated gene.

### 3) Differential display:

This technique is described by Chuang *et al.*, *J. Bacteriol.* 175:2026-2036 (1993), the contents of which is incorporated by reference for background purposes. This method identifies those genes which are expressed in an organism by identifying mRNA present using randomly-primed RT-PCR. By comparing pre-infection and post infection profiles, genes up and down regulated during infection can be identified and the RT-PCR product sequenced and matched to library sequences.

### 4) Generation of conditional lethal mutants by transposon mutagenesis:

This technique, described by de Lorenzo, V. *et al.*, *Gene* 123:17-24 (1993); Neuwald, A. F. *et al.*, *Gene* 125: 69-73 (1993); and Takiff, H. E. *et al.*, *J. Bacteriol.* 174:1544-1553(1992), the contents of which is incorporated by reference for background purposes, identifies genes whose expression are essential for cell viability.

In this technique transposons carrying controllable promoters, which provide transcription outward from the transposon in one or both directions, are generated. Random insertion of these transposons into target organisms and subsequent isolation of insertion mutants in the presence of inducer of promoter activity ensures that insertions which separate promoter from coding region of a gene whose expression is essential for cell viability will be recovered. Subsequent replica plating in the absence of inducer identifies such insertions, since they fail to survive. Sequencing of the flanking regions of the transposon allows identification of site of insertion and identification of the gene disrupted. Close monitoring of the changes in cellular processes/morphology during growth in the absence of inducer yields information on likely function of the gene. Such monitoring could include flow cytometry (cell division, lysis, redox potential, DNA replication), incorporation of radiochemically labeled precursors into DNA, RNA, protein, lipid, peptidoglycan, monitoring reporter enzyme gene fusions which respond to known cellular stresses.

### 5) Generation of conditional lethal mutants by chemical mutagenesis:

This technique is described by Beckwith, J., *Methods in Enzymology* 204: 3-18(1991), the contents of which are incorporated herein by reference for background purposes. In this technique random chemical mutagenesis of target organism, growth at temperature other than physiological temperature (permissive temperature) and subsequent replica plating and growth at different temperature (e.g. 42°C to identify ts, 25°C to identify cs) are used to identify those isolates which now fail to grow (conditional mutants). As above close monitoring of the changes upon growth at the non-permissive temperature yields information on the function of the mutated gene. Complementation of conditional lethal mutation by library from target organism and sequencing of complementing gene allows matching with library sequences.

Each of these techniques may have advantages or disadvantage depending on the particular application. The skilled artisan would choose the approach that is the most relevant with the particular end use in mind. For example, some genes might be recognised as essential for infection but in reality are only necessary for the initiation of infection and so their products would represent relatively unattractive targets for antibacterials developed to cure established and chronic infections.

### 6) RT-PCR:

*Streptococcus pneumoniae* messenger RNA is isolated from bacterial infected tissue e.g. 48 hour murine lung infections, and the amount of each mRNA species assessed by reverse transcription of the RNA sample primed with random hexanucleotides followed by PCR with gene specific primer pairs. The determination of the presence and amount of a particular mRNA species by quantification of the resultant PCR product provides information on the bacterial genes which are transcribed in the infected tissue. Analysis of gene transcription can be carried out at different times of infection to gain a detailed knowledge of gene regulation in bacterial pathogenesis allowing for a clearer understanding of which gene products represent targets for screens for novel antibacterials. Because of the gene specific nature of the PCR primers employed it should be understood that the bacterial mRNA preparation need not be free of mammalian RNA. This allows the investigator to carry out a simple and quick RNA preparation from infected tissue to obtain bacterial mRNA species which are very short lived in the bacterium (in the order of 2 minute half-lives).



Optimally the bacterial mRNA is prepared from infected murine lung tissue by mechanical disruption in the presence of TRIzol (GIBCO-BRL) for very short periods of time, subsequent processing according to the manufacturers of TRIzol reagent and DNAase treatment to remove contaminating DNA. Preferably the process is optimised by finding those conditions which give a maximum amount of *S. pneumoniae* 16S ribosomal RNA as detected by probing Northern blots with a suitably labelled sequence specific oligonucleotide probe. Typically a 5' dye labelled primer is used in each PCR primer pair in a PCR reaction which is terminated optimally between 8 and 25 cycles. The PCR products are separated on 6% polyacrylamide gels with detection and quantification using GeneScanner (manufactured by ABI).

Use of the of these technologies when applied to the sequences of the present invention enables identification of bacterial proteins expressed during infection, inhibitors of which would have utility in anti-bacterial therapy.

## Polynucleotides

In accordance with one aspect of the present invention, there are provided isolated polynucleotides which encode the IgAFcBP polypeptide having the deduced amino acid sequence of Figure 2 [SEQ ID NO:2].

Using the information provided herein, such as the polynucleotide sequence set out in Figure 1 [SEQ ID NO: 1], a polynucleotide of the present invention encoding IgAFcBP polypeptide may be obtained using standard cloning and screening procedures, such as those for cloning and sequencing chromosomal DNA fragments from *Streptococcus pneumoniae* 0100993 cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide of the invention sequence, such as that sequence given in Figure 1 [SEQ ID NO: 1] typically a library of clones of chromosomal DNA of *Streptococcus pneumoniae* 0100993 in *E. coli* or some other suitable host is probed with a radiolabelled oligonucleotide, preferably a 17mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using high stringency washes. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently such sequencing is performed using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Illustrative of the invention, the polynucleotide set out in Figure 1 [SEQ ID NO:1] was discovered in a DNA library derived from *Streptococcus pneumoniae* 0100993.

IgAFcBP of the invention is structurally related to other streptococcal IgAFcBPs, as shown by the results of sequencing the cDNA encoding IgAFcBP in the deposited clone. The DNA sequence thus obtained is set out in Figure 1 [SEQ ID NO: 1]. It contains an open reading frame encoding a protein of about 170 amino acid residues. The protein exhibits greatest homology to streptococcal IgA Fc binding protein among known proteins. IgAFcBP of Figure 2 [SEQ ID NO:2] has about 50% identity and about 60% similarity with the amino acid sequence (amino acid residues 722-822) of the IgA Fc binding protein of *Streptococcus agalactiae* (SWISSPROT:BAG\_STRAG).

Polynucleotides of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The DNA may be double-stranded or single-stranded. Single-stranded DNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

The coding sequence which encodes the polypeptide may be identical to the coding sequence of the polynucleotide shown in Figure 1 [SEQ ID NO: 1]. It also may be a polynucleotide with a different sequence, which, as a result of the redundancy (degeneracy) of the genetic code, encodes the polypeptide of the DNA of Figure 1 [SEQ ID NO: 1].

Polynucleotides of the present invention which encode the polypeptide of Figure 2 [SEQ ID NO:2] may include, but are not limited to the coding sequence for the mature polypeptide, by itself; the coding sequence for the mature polypeptide and additional coding sequences, such as those encoding a leader or secretory sequence, such as a pre-, or pro- or prepro- protein sequence; the coding sequence of the mature polypeptide, with or without the aforementioned additional coding sequences, together with additional, non-coding sequences, including for example, but not limited to non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription (including termination signals, for example), ribosome binding, mRNA stability elements, and additional coding sequence which encode additional amino acids, such as those which provide additional functionalities. Thus, for instance, the polypeptide may be fused to a marker sequence, such as a peptide, which facilitates purification of the fused polypeptide. In certain embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, such as the tag provided in the pQE vector (Qiagen, Inc.), among others, many of which are commercially available. As described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. The HA tag may also be used to create fusion proteins and corresponds to an epitope derived of influenza hemagglutinin protein, which has been described by Wilson *et al.*, *Cell* 37: 767 (1984), for instance.

In accordance with the foregoing, the term "polynucleotide encoding a polypeptide" as used herein encompasses

polynucleotides which include a sequence encoding a polypeptide of the present invention, particularly bacterial, and more particularly streptococcal IgAFcBP having the amino acid sequence set out in Figure 2 [SEQ ID NO:2]. The term encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, interrupted by integrated phage or insertion sequence or editing) together with additional regions, that also may contain coding and/or non-coding sequences.

The present invention further relates to variants of the herein above described polynucleotides which encode for fragments, analogs and derivatives of the polypeptide having the deduced amino acid sequence of Figure 2 [SEQ ID NO:2]. A variant of the polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant that is not known to occur naturally. Such non-naturally occurring variants of the polynucleotide may be made by mutagenesis techniques, including those applied to polynucleotides, cells or organisms.

Among variants in this regard are variants that differ from the aforementioned polynucleotides by nucleotide substitutions, deletions or additions. The substitutions, deletions or additions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions.

Among the particularly preferred embodiments of the invention in this regard are polynucleotides encoding polypeptides having the amino acid sequence of IgAFcBP set out in Figure 2 [SEQ ID NO:2]; variants, analogs, derivatives and fragments thereof, and fragments of the variants, analogs and derivatives.

Further particularly preferred in this regard are polynucleotides encoding IgAFcBP variants, analogs, derivatives and fragments, and variants, analogs and derivatives of the fragments, which have the amino acid sequence of IgAFcBP polypeptide of Figure 2 [SEQ ID NO:2] in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of IgAFcBP. Also especially preferred in this regard are conservative substitutions. Most highly preferred are polynucleotides encoding polypeptides having the amino acid sequence of Figure 2 [SEQ ID NO:2], without substitutions.

Further preferred embodiments of the invention are polynucleotides that are at least 70% identical to a polynucleotide encoding IgAFcBP polypeptide having the amino acid sequence set out in Figure 2 [SEQ ID NO:2], and polynucleotides which are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 80% identical to a polynucleotide encoding IgAFcBP polypeptide of the *Streptococcus pneumoniae* DNA of the deposited clone and polynucleotides complementary thereto. In this regard, polynucleotides at least 90% identical to the same are particularly preferred, and among these particularly preferred polynucleotides, those with at least 95% are especially preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Particularly preferred embodiments in this respect, moreover, are polynucleotides which encode polypeptides which retain substantially the same biological function or activity as the mature polypeptide encoded by the DNA of Figure 1 [SEQ ID NO:1].

The present invention further relates to polynucleotides that hybridize to the herein above-described sequences. In this regard, the present invention especially relates to polynucleotides which hybridize under stringent conditions to the herein above-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences.

As discussed additionally herein regarding polynucleotide assays of the invention, for instance, polynucleotides of the invention as discussed above, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding IgAFcBP and to isolate cDNA and genomic clones of other genes that have a high sequence similarity to the IgAFcBP gene. Such probes generally will comprise at least 15 bases. Preferably, such probes will have at least 30 bases and may have at least 50 bases. Particularly preferred probes will have at least 30 bases and will have 50 bases or less.

For example, the coding region of the IgAFcBP gene may be isolated by screening using the known DNA sequence to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the present invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

The polynucleotides and polypeptides of the present invention may be employed as research reagents and materials for discovery of treatments of and diagnostics for disease, particularly human disease, as further discussed herein relating to polynucleotide assays, *inter alia*.

The polynucleotides of the invention that are oligonucleotides, including sequences that may be used as PCR primers in the process herein described to determine whether or not the *Streptococcus pneumoniae* genes identified herein in whole or in part are transcribed in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The polynucleotides may encode a polypeptide which is the mature protein plus additional amino or carboxyl-

terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

A precursor protein, having the mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

In sum, a polynucleotide of the present invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences which are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

#### Deposited materials

The deposited material is a cloned *Streptococcus pneumoniae*, strain 0100993 has been deposited at the National Collection of Industrial and Marine Bacteria Ltd. (NCIMB), Aberdeen, Scotland under number NCIMB 40794 on 11 April 1996. Also as noted above, the *Streptococcus pneumoniae* deposit and polynucleotides derived therefrom using methods known in the art or described herein are referred to herein as "the deposited clone" or as "the DNA of the deposited clone."

The deposit has been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent Procedure. The strain will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposit is provided merely as convenience to those of skill in the art and is not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

The sequence of the polynucleotides contained in the deposited material, as well as the amino acid sequence of the polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

A license may be required to make, use or sell the deposited materials, and no such license is hereby granted.

#### Polypeptides

The present invention further relates to a prokaryotic IgAFcBP polypeptide which has the deduced amino acid sequence of Figure 2 [SEQ ID NO:2].

The invention also relates to fragments, analogs and derivatives of these polypeptides. The terms "fragment," "derivative" and "analog" when referring to the polypeptide of Figure 2 [SEQ ID NO:2], means a polypeptide which retains essentially the same biological function or activity as such polypeptide. Thus, an analog includes a proprotein which can be activated by cleavage of the proprotein portion to produce an active mature polypeptide.

The polypeptide of the present invention may be a recombinant polypeptide, a natural polypeptide or a synthetic polypeptide. In certain preferred embodiments it is a recombinant polypeptide.

The fragment, derivative or analog of the polypeptide of Figure 2 [SEQ ID NO:2] may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

Among the particularly preferred embodiments of the invention in this regard are polypeptides having the amino acid sequence of IgAFcBP set out in Figure 2 [SEQ ID NO:2], variants, analogs, derivatives and fragments thereof, and variants, analogs and derivatives of the fragments. Alternatively, particularly preferred embodiments of the invention in this regard are polypeptides having the amino acid sequence of the IgAFcBP, variants, analogs, derivatives and fragments thereof, and variants, analogs and derivatives of the fragments.

Among preferred variants are those that vary from a reference by conservative amino acid substitutions. Such substitutions are those that substitute a given amino acid in a polypeptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu and Ile; interchange of the hydroxyl residues Ser and Thr, exchange of the acidic residues Asp and Glu, substitution between the amide residues Asn and Gln, exchange of the basic residues Lys and Arg and replacements

among the aromatic residues Phe, Tyr.

Further particularly preferred in this regard are variants, analogs, derivatives and fragments, and variants, analogs and derivatives of the fragments, having the amino acid sequence of the IgAFcBP polypeptide of Figure 2 [SEQ ID NO:2], in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, deleted or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the IgAFcBP. Also especially preferred in this regard are conservative substitutions. Most highly preferred are polypeptides having the amino acid sequence of Figure 2 [SEQ ID NO:2] without substitutions.

The polypeptides and polynucleotides of the present invention are preferably provided in an isolated form, and preferably are purified to homogeneity.

The polypeptides of the present invention include the polypeptide of Figure 2 [SEQ ID NO:2] (in particular the mature polypeptide) as well as polypeptides which have at least 80% identity to the polypeptide of Figure 2 [SEQ ID NO:2] and more preferably at least 90% similarity (more preferably at least 90% identity) to the polypeptide of Figure 2 [SEQ ID NO:2] and still more preferably at least 95% similarity (still more preferably at least 95% identity) to the polypeptide of Figure 2 [SEQ ID NO:2] and also include portions of such polypeptides with such portion of the polypeptide generally containing at least 30 amino acids and more preferably at least 50 amino acids.

Fragments or portions of the polypeptides of the present invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, the fragments may be employed as intermediates for producing the full-length polypeptides. Fragments or portions of the polynucleotides of the present invention may be used to synthesize full-length polynucleotides of the present invention.

## ^ Fragments

Also among preferred embodiments of this aspect of the present invention are polypeptides comprising fragments of IgAFcBP, most particularly fragments of IgAFcBP having the amino acid set out in Figure 2 [SEQ ID NO:2], and fragments of variants and derivatives of the IgAFcBP of Figure 2 [SEQ ID NO:2].

In this regard a fragment is a polypeptide having an amino acid sequence that entirely is the same as part but not all of the amino acid sequence of the aforementioned IgAFcBP polypeptides and variants or derivatives thereof.

Such fragments may be "free-standing," *i.e.*, not part of or fused to other amino acids or polypeptides, or they may be comprised within a larger polypeptide of which they form a part or region. When comprised within a larger polypeptide, the presently discussed fragments most preferably form a single continuous region. However, several fragments may be comprised within a single larger polypeptide. For instance, certain preferred embodiments relate to a fragment of a IgAFcBP polypeptide of the present comprised within a precursor polypeptide designed for expression in a host and having heterologous pre and pro-polypeptide regions fused to the amino terminus of the IgAFcBP fragment and an additional region fused to the carboxyl terminus of the fragment. Therefore, fragments in one aspect of the meaning intended herein, refers to the portion or portions of a fusion polypeptide or fusion protein derived from IgAFcBP.

As representative examples of polypeptide fragments of the invention, there may be mentioned those which have from about 5-15, 10-20, 15-40, 30-55, 41-75, 41-80, 41-90, 50-100, 75-100, 90-115, and 100-125 amino acids long. Particular examples of polypeptide fragments of the inventions that may be mentioned include fragments from amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, and 101-120, 121-140, 141-160, and 161-170, and any combination of these 20mer fragments.

In this context about includes the particularly recited range and ranges larger or smaller by several, a few, 5, 4, 3, 2 or 1 amino acid at either extreme or at both extremes. For instance, about 40-90 amino acids in this context means a polypeptide fragment of 40 plus or minus several, a few, 5, 4, 3, 2 or 1 amino acids to 90 plus or minus several a few, 5, 4, 3, 2 or 1 amino acid residues, *i.e.*, ranges as broad as 40 minus several amino acids to 90 plus several amino acids to as narrow as 40 plus several amino acids to 90 minus several amino acids.

Highly preferred in this regard are the recited ranges plus or minus as many as 5 amino acids at either or at both extremes. Particularly highly preferred are the recited ranges plus or minus as many as 3 amino acids at either or at both the recited extremes. Especially particularly highly preferred are ranges plus or minus 1 amino acid at either or at both extremes or the recited ranges with no additions or deletions.

Among especially preferred fragments of the invention are truncation mutants of IgAFcBP. Truncation mutants include IgAFcBP polypeptides having the amino acid sequence of Figure 2, or of variants or derivatives thereof, except for deletion of a continuous series of residues (that is, a continuous region, part or portion) that includes the amino terminus, or a continuous series of residues that includes the carboxyl terminus or, as in double truncation mutants, deletion of two continuous series of residues, one including the amino terminus and one including the carboxyl terminus. Fragments having the size ranges set out about also are preferred embodiments of truncation fragments, which are especially preferred among fragments generally. Degradation forms of the polypeptides of the invention in a host cell, particularly a mammal, and especially a human, are also preferred. Also preferred is a fragment of polypeptide expressed from the polynucleotide construct used for during genetic immunization, for example, small peptide fragments

formed by during secretion and presentation on the MHComplex.

Also preferred in this aspect of the invention are fragments characterized by structural or functional attributes of IgAFcBP. Preferred embodiments of the invention in this regard include fragments that comprise alpha-helix and alpha-helix forming regions ("alpha-regions"), beta-sheet and beta-sheet-forming regions ("beta-regions"), turn and turn-forming regions ("turn-regions"), coil and coil-forming regions ("coil-regions"), hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions and high antigenic index regions of IgAFcBP.

Among highly preferred fragments in this regard are those that comprise regions of IgAFcBP that combine several structural features, such as several of the features set out above. In this regard, the regions defined by the residues about 10 to about 20, about 40 to about 50, about 70 to about 90 and about 100 to about 170 of Figure 2 [SEQ ID NO: 2], which all are characterized by amino acid compositions highly characteristic of turn-regions, hydrophilic regions, flexible-regions, surface-forming regions, and high antigenic index-regions, are especially highly preferred regions. Such regions may be comprised within a larger polypeptide or may be by themselves a preferred fragment of the present invention, as discussed above. It will be appreciated that the term "about" as used in this paragraph has the meaning set out above regarding fragments in general.

Further preferred regions are those that mediate activities of IgAFcBP. Most highly preferred in this regard are fragments that have a chemical, biological or other activity of IgAFcBP, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Highly preferred in this regard are fragments that contain regions that are homologs in sequence, or in position, or in both sequence and to active regions of related polypeptides, such as the related polypeptides set out in Figure 2 [SEQ ID NO:2], which include streptococcal IgAFcBP. Among particularly preferred fragments in these regards are truncation mutants, as discussed above. Further preferred polynucleotide fragments are those that are antigenic or immunogenic in an animal, especially in a human.

It will be appreciated that the invention also relates to, among others, polynucleotides encoding the aforementioned fragments, polynucleotides that hybridize to polynucleotides encoding the fragments, particularly those that hybridize under stringent conditions, and polynucleotides, such as PCR primers, for amplifying polynucleotides that encode the fragments. In these regards, preferred polynucleotides are those that correspond to the preferred fragments, as discussed above.

#### Vectors, host cells, expression

The present invention also relates to vectors which comprise a polynucleotide or polynucleotides of the present invention, host cells which are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques.

Host cells can be genetically engineered to incorporate polynucleotides and express polypeptides of the present invention. For instance, polynucleotides may be introduced into host cells using well known techniques of infection, transduction, transfection, transvection and transformation. The polynucleotides may be introduced alone or with other polynucleotides. Such other polynucleotides may be introduced independently, co-introduced or introduced joined to the polynucleotides of the invention.

Thus, for instance, polynucleotides of the invention may be transfected into host cells with another, separate, polynucleotide encoding a selectable marker, using standard techniques for co-transfection and selection in, for instance, mammalian cells. In this case the polynucleotides generally will be stably incorporated into the host cell genome.

Alternatively, the polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. The vector construct may be introduced into host cells by the aforementioned techniques. Generally, a plasmid vector is introduced as DNA in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. Electroporation also may be used to introduce polynucleotides into a host. If the vector is a virus, it may be packaged in vitro or introduced into a packaging cell and the packaged virus may be transduced into cells. A wide variety of techniques suitable for making polynucleotides and for introducing polynucleotides into cells in accordance with this aspect of the invention are well known and routine to those of skill in the art. Such techniques are reviewed at length in et al. cited above, which is illustrative of the many laboratory manuals that detail these techniques. In accordance with this aspect of the invention the vector may be, for example, a plasmid vector, a single or double-stranded phage vector, a single or double-stranded RNA or DNA viral vector. Such vectors may be introduced into cells as polynucleotides, preferably DNA, by well known techniques for introducing DNA and RNA into cells. The vectors, in the case of phage and viral vectors also may be and preferably are introduced into cells as packaged or encapsidated virus by well known techniques for infection and transduction. Viral vectors may be replication competent or replication defective. In the latter case viral propagation generally will occur only in complementing host cells.

Preferred among vectors, in certain respects, are those for expression of polynucleotides and polypeptides of the present invention. Generally, such vectors comprise cis-acting control regions effective for expression in a host operatively linked to the polynucleotide to be expressed. Appropriate trans-acting factors either are supplied by the host,

supplied by a complementing vector or supplied by the vector itself upon introduction into the host.

In certain preferred embodiments in this regard, the vectors provide for specific expression. Such specific expression may be inducible expression or expression only in certain types of cells or both inducible and cell-specific. Particularly preferred among inducible vectors are vectors that can be induced for expression by environmental factors that are easy to manipulate, such as temperature and nutrient additives. A variety of vectors suitable to this aspect of the invention, including constitutive and inducible expression vectors for use in prokaryotic and eukaryotic hosts, are well known and employed routinely by those of skill in the art.

The engineered host cells can be cultured in conventional nutrient media, which may be modified as appropriate for, *inter alia*, activating promoters, selecting transformants or amplifying genes. Culture conditions, such as temperature, pH and the like, previously used with the host cell selected for expression generally will be suitable for expression of polypeptides of the present invention as will be apparent to those of skill in the art.

A great variety of expression vectors can be used to express a polypeptide of the invention. Such vectors include, among others, chromosomal, episomal and virus-derived vectors, *e.g.*, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids, all may be used for expression in accordance with this aspect of the present invention. Generally, any vector suitable to maintain, propagate or express polynucleotides to express a polypeptide in a host may be used for expression in this regard.

The appropriate DNA sequence may be inserted into the vector by any of a variety of well-known and routine techniques. In general, a DNA sequence for expression is joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more restriction endonucleases and then joining the restriction fragments together using T4 DNA ligase. Procedures for restriction and ligation that can be used to this end are well known and routine to those of skill. Suitable procedures in this regard, and for constructing expression vectors using alternative techniques, which also are well known and routine to those of skill, are set forth in great detail in Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989).

The DNA sequence in the expression vector is operatively linked to appropriate expression control sequence(s), including, for instance, a promoter to direct mRNA transcription. Representatives of such promoters include the phage lambda PL promoter, the *E. coli* lac, trp and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name just a few of the well-known promoters. It will be understood that numerous promoters not mentioned are suitable for use in this aspect of the invention are well known and readily may be employed by those of skill in the manner illustrated by the discussion and the examples herein.

In general, expression constructs will contain sites for transcription initiation and termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the constructs will include a translation initiating AUG at the beginning and a termination codon appropriately positioned at the end of the polypeptide to be translated.

In addition, the constructs may contain control regions that regulate as well as engender expression. Generally, in accordance with many commonly practiced procedures, such regions will operate by controlling transcription, such as transcription factors, repressor binding sites and termination, among others.

Vectors for propagation and expression generally will include selectable markers. Such markers also may be suitable for amplification or the vectors may contain additional markers for this purpose. In this regard, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells. Preferred markers include dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, and tetracycline or ampicillin resistance genes for culturing *E. coli* and other prokaryotes.

The vector containing the appropriate DNA sequence as described elsewhere herein, as well as an appropriate promoter, and other appropriate control sequences, may be introduced into an appropriate host using a variety of well known techniques suitable to expression therein of a desired polypeptide. Representative examples of appropriate hosts include bacterial cells, such as streptococci, staphylococci, *E. coli*, streptomyces and *Salmonella typhimurium* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS and Bowes melanoma cells; and plant cells. Hosts for of a great variety of expression constructs are well known, and those of skill will be enabled by the present disclosure readily to select a host for expressing a polypeptides in accordance with this aspect of the present invention.

More particularly, the present invention also includes recombinant constructs, such as expression constructs, comprising one or more of the sequences described above. The constructs comprise a vector, such as a plasmid or viral vector, into which such a sequence of the invention has been inserted. The sequence may be inserted in a forward or reverse orientation. In certain preferred embodiments in this regard, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and



promoters are known to those of skill in the art, and there are many commercially available vectors suitable for use in the present invention.

The following vectors, which are commercially available, are provided by way of example. Among vectors preferred for use in bacteria are pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. These vectors are listed solely by way of illustration of the many commercially available and well known vectors that are available to those of skill in the art for use in accordance with this aspect of the present invention. It will be appreciated that any other plasmid or vector suitable for, for example, introduction, maintenance, propagation or expression of a polynucleotide or polypeptide of the invention in a host may be used in this aspect of the invention.

Promoter regions can be selected from any desired gene using vectors that contain a reporter transcription unit lacking a promoter region, such as a chloramphenicol acetyl transferase ("CAT") transcription unit, downstream of restriction site or sites for introducing a candidate promoter fragment; *i.e.*, a fragment that may contain a promoter. As is well known, introduction into the vector of a promoter-containing fragment at the restriction site upstream of the *cat* gene engenders production of CAT activity, which can be detected by standard CAT assays. Vectors suitable to this end are well known and readily available. Two such vectors are pKK232-8 and pCM7. Thus, promoters for expression of polynucleotides of the present invention include not only well known and readily available promoters, but also promoters that readily may be obtained by the foregoing technique, using a reporter gene.

Among known prokaryotic promoters suitable for expression of polynucleotides and polypeptides in accordance with the present invention are the *E. coli* *lacI* and *lacZ* and promoters, the T3 and T7 promoters, the *gpt* promoter, the lambda PR, PL promoters and the *trp* promoter.

Among known eukaryotic promoters suitable in this regard are the CMV immediate early promoter, the HSV thymidine kinase promoter, the early and late SV40 promoters, the promoters of retroviral LTRs, such as those of the Rous sarcoma virus ("RSV"), and metallothionein promoters, such as the mouse metallothionein-I promoter.

Selection of appropriate vectors and promoters for expression in a host cell is a well known procedure and the requisite techniques for expression vector construction, introduction of the vector into the host and expression in the host are routine skills in the art.

The present invention also relates to host cells containing the above-described constructs discussed above. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell.

Introduction of the construct into the host cell can be affected by calcium phosphate transfection, DEAE-dextran mediated transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

Constructs in host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence. Alternatively, the polypeptides of the invention can be synthetically produced by conventional peptide synthesizers.

Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

Generally, recombinant expression vectors will include origins of replication, a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence, and a selectable marker to permit isolation of vector containing cells after exposure to the vector. Among suitable promoters are those derived from the genes that encode glycolytic enzymes such as 3-phosphoglycerate kinase ("PGK"),  $\alpha$ -factor, acid phosphatase, and heat shock proteins, among others. Selectable markers include the ampicillin resistance gene of *E. coli* and the *trp1* gene of *S. cerevisiae*.

Transcription of the DNA encoding the polypeptides of the present invention by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp that act to increase transcriptional activity of a promoter in a given host cell-type. Examples of enhancers include the SV40 enhancer, which is located on the late side of the replication origin at bp 100 to 270, the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers:

Polynucleotides of the invention, encoding the heterologous structural sequence of a polypeptide of the invention generally will be inserted into the vector using standard techniques so that it is operably linked to the promoter for



expression. The polynucleotide will be positioned so that the transcription start site is located appropriately 5' to a ribosome binding site. The ribosome binding site will be 5' to the AUG that initiates translation of the polypeptide to be expressed. Generally, there will be no other open reading frames that begin with an initiation codon, usually AUG, and lie between the ribosome binding site and the initiation codon. Also, generally, there will be a translation stop codon at the end of the polypeptide and there will be a polyadenylation signal in constructs for use in eukaryotic hosts. Transcription termination signal appropriately disposed at the 3' end of the transcribed region may also be included in the polynucleotide construct.

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals but also additional heterologous functional regions. Thus, for instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence in the host cell, during purification or during subsequent handling and storage. Also, region also may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability or to facilitate purification, among others, are familiar and routine techniques in the art. A preferred fusion protein comprises a heterologous region from immunoglobulin that is useful to solubilize or purify polypeptides. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in fusion protein is thoroughly advantageous for use in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (see EP-A 0232 262). On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified in the advantageous manner described. This is the case when Fc portion proves to be a hindrance to use in assays, therapy or diagnosis, for example when the fusion protein is to be used as antigen for immunizations. In drug discovery, for example, human proteins, such as, shIL5- has been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett et al., *Journal of Molecular Recognition*, Vol. 8 52-58 (1995) and K. Johanson et al., *The Journal of Biological Chemistry*, Vol. 270, No. 16, pp 9459-9471 (1995).

Suitable prokaryotic hosts for propagation, maintenance or expression of polynucleotides and polypeptides in accordance with the invention include streptococci, *Escherichia coli*, *Bacillus subtilis* and *Salmonella typhimurium*. Various species of *Pseudomonas*, *Streptomyces*, and *Staphylococcus* are also suitable hosts in this regard. Moreover, many other hosts also known to those of skill may be employed in this regard.

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed.

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, where the selected promoter is inducible it is induced by appropriate means (e.g., temperature shift or exposure to chemical inducer) and cells are cultured for an additional period.

Cells typically then are harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, such methods are well known to those skilled in the art.

Various mammalian cell culture systems can be employed for expression, as well. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblast, described in Gluzman et al., *Cell* 23: 175 (1981). Other cell lines capable of expressing a compatible vector include for example, the C127, 3T3, CHO, HeLa, human kidney 293 and BHK cell lines.

Mammalian expression vectors may comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation sites, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking non-transcribed sequences that are necessary for expression. In certain preferred embodiments in this regard DNA sequences derived from the SV40 splice sites, and the SV40 polyadenylation sites are used for required non-transcribed genetic elements of these types.

IgAFcBP polypeptide can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for

purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and or purification.

Polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

IgAFcBP polynucleotides and polypeptides may be used in accordance with the present invention for a variety of applications, particularly those that make use of the chemical and biological properties of IgAFcBP. Additional applications relate to diagnosis and to treatment of disorders of cells, tissues and organisms. These aspects of the invention are illustrated further by the following discussion.

#### Polynucleotide assays

This invention is also related to the use of the IgAFcBP polynucleotides to detect complementary polynucleotides such as, for example, as a diagnostic reagent. Detection of IgAFcBP in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method that can add to, define or allow a diagnosis of a disease. Eukaryotes (herein also "individual(s)"), particularly mammals, and especially humans, carrying a IgAFcBP gene may be detected at the DNA level by a variety of techniques. For example, individuals carrying a IgAFcBP gene may be detected at the DNA level by a variety of techniques. Nucleic acids for diagnosis may be obtained from an individual's cells and tissues, such as bone, blood, muscle, cartilage, and skin. Tissue biopsy and autopsy material is also preferred for samples from an individual to use in a diagnostic assay. Genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR prior to analysis. PCR (Saiki et al., *Nature*, 324: 163-166 (1986)). RNA or cDNA may also be used in the same ways. As an example, PCR primers complementary to the nucleic acid encoding IgAFcBP can be used to identify and analyze IgAFcBP presence and expression. Using PCR, characterization of the strain of prokaryote present in a eukaryote, particularly a mammal, and especially a human, may be made by an analysis of the genotype of the prokaryote gene. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the genotype of a reference sequence. Point mutations can be identified by hybridizing amplified DNA to radiolabeled IgAFcBP RNA or alternatively, radiolabeled IgAFcBP antisense DNA sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

Sequence differences between a reference gene and genes having mutations also may be revealed by direct DNA sequencing. In addition, cloned DNA segments may be employed as probes to detect specific DNA segments. The sensitivity of such methods can be greatly enhanced by appropriate use of PCR or another amplification method. For example, a sequencing primer is used with double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence determination is performed by conventional procedures with radiolabeled nucleotide or by automatic sequencing procedures with fluorescent-tags.

Genetic testing based on DNA sequence differences may be achieved by detection of alteration in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures (see, e.g., Myers et al., *Science*, 230: 1242 (1985)).

Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (e.g., Cotton et al., *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985)).

Thus, the detection of a specific DNA sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing or the use of restriction enzymes, (e.g., restriction fragment length polymorphisms ("RFLP") and Southern blotting of genomic DNA.

In addition to more conventional gel-electrophoresis and DNA sequencing, mutations also can be detected by *in situ* analysis.

A mutation may be ascertained for example, by a DNA sequencing assay. Samples are processed by methods known in the art to capture the RNA. First strand cDNA is synthesized from the RNA samples by adding an oligonucleotide primer consisting of sequences which hybridize to a region on the mRNA. Reverse transcriptase and deoxynucleotides are added to allow synthesis of the first strand cDNA. Primer sequences are synthesized based on the DNA sequence of the IgAFcBP protein of the invention. The primer sequence is generally comprised of at least 15 consecutive bases, and may contain at least 30 or even 50 consecutive bases.

The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR (Saiki et

*al.*, *Nature*, 324:163-166 (1986)) prior to analysis. RT-PCR can also be used to detect mutations. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. R or cDNA may also be used for the same purpose, PCR or RT-PCR. As an example, PCR primers complementary to the nucleic acid encoding IgAFcBP can be used to identify and analyze mutations. Examples of representative primers are shown below in Table 1. For example, deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to diolabeled RNA or alternatively, radiolabeled antisense DNA sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase A digestion or by differences in melting temperatures.

Table 1

## Primers used for detection of IgAFcBP gene

SEQ ID NO:

1. 5' ATTAAGCGAA CGTATTTGTA 3'

2. 5' GGAATTTAGT CACAACGTGCA 3'

The above primers may be used for amplifying IgAFcBP DNA isolated from a sample derived from an individual. The invention also provides the primers of Table 1 with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. The primers may be used to amplify the gene isolated from the individual such that the gene may then be subjected to various techniques for elucidation of the DNA sequence. In this way, mutations in the DNA sequence may be diagnosed.

Sequence differences between the reference gene and genes having mutations may be revealed by the direct DNA sequencing method. In addition, cloned DNA segments may be employed as probes to detect specific DNA segments. The sensitivity of this method is greatly enhanced when combined with PCR. For example, a sequencing primer is used with double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence determination is performed by conventional procedures with radiolabeled nucleotide or by automatic sequencing procedures with fluorescent-tags.

Genetic testing based on DNA sequence differences may be achieved by detection of alteration in electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Small sequence deletions and insertions can be visualized by high resolution gel electrophoresis. DNA fragments of different sequences may be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures (see, e.g., Myers *et al.*, *Science*, 230:122 (1985)).

Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase S1 protection or the chemical cleavage method (e.g., Cotton *et al.*, *PNAS*, USA, 85:4397-4401 (1988)).

Thus, the detection of a specific DNA sequence and/or quantitation of the level of the sequence may be achieved by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing or the use of restriction enzymes, (e.g., Restriction Fragment Length Polymorphisms (RFLP)) and Southern blotting of genomic DNA.

The invention provides a process for diagnosing, disease, particularly otitis media, conjunctivitis, pneumonia, bacteremia, meningitis, sinusitis, pleural empyema and endocarditis, and most particularly meningitis, such as for example infection of cerebrospinal fluid, comprising determining from a sample derived from an individual a increased level of expression of polynucleotide having the sequence of Figure 1 [SEQ ID NO: 1]. Increased expression of polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

In addition to more conventional gel-electrophoresis and DNA sequencing, mutations can also be detected by *situ* analysis.

**Polypeptide assays**

The present invention also relates to a diagnostic assays such as quantitative and diagnostic assays for detecting levels of IgAFcBP protein in cells and tissues, including determination of normal and abnormal levels. Thus, for instance a diagnostic assay in accordance with the invention for detecting over-expression of IgAFcBP protein compared

normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a protein, such as an IgAFcBP protein of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays. Among these ELISAs frequently are preferred. An ELISA assay initially comprises preparing an antibody specific to IgAFcBP, preferably a monoclonal antibody. In addition a reporter antibody generally is prepared which binds to the monoclonal antibody. The reporter antibody is attached a detectable reagent such as radioactive, fluorescent or enzymatic reagent, in this example horseradish peroxidase enzyme.

To carry out an ELISA a sample is removed from a host and incubated on a solid support, e.g., a polystyrene dish, that binds the proteins in the sample. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein such as bovine serum albumin. Next, the monoclonal antibody is incubated in the dish during which time the monoclonal antibodies attach to any IgAFcBP proteins attached to the polystyrene dish. Unbound monoclonal antibody is washed out with buffer. The reporter antibody linked to horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to IgAFcBP. Unattached reporter antibody is then washed out. Reagents for peroxidase activity, including a colorimetric substrate are then added to the dish. Immobilized peroxidase, linked to IgAFcBP through the primary and secondary antibodies, produces a colored reaction product. The amount of color developed in a given time period indicates the amount of IgAFcBP protein present in the sample. Quantitative results typically are obtained by reference to a standard curve.

A competition assay may be employed wherein antibodies specific to IgAFcBP attached to a solid support and labeled IgAFcBP and a sample derived from the host are passed over the solid support and the amount of label detected attached to the solid support can be correlated to a quantity of IgAFcBP in the sample.

### Antibodies

The polypeptides, their fragments or other derivatives, or analogs thereof, or cells expressing them can be used as an immunogen to produce antibodies thereto. These antibodies can be, for example, polyclonal or monoclonal antibodies. The present invention also includes chimeric, single chain, and humanized antibodies, as well as Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

Antibodies generated against the polypeptides corresponding to a sequence of the present invention can be obtained by direct injection of the polypeptides into an animal or by administering the polypeptides to an animal, preferably a nonhuman. The antibody so obtained will then bind the polypeptides itself. In this manner, even a sequence encoding only a fragment of the polypeptides can be used to generate antibodies binding the whole native polypeptides. Such antibodies can then be used to isolate the polypeptide from tissue expressing that polypeptide.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* 4: 72 (1983) and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies to immunogenic polypeptide products of this invention.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or purify the polypeptide of the present invention by attachment of the antibody to a solid support for isolation and/or purification by affinity chromatography.

Thus, among others, antibodies against IgAFcBP may be employed to inhibit or treat infections, particularly bacterial infections and especially otitis media, conjunctivitis, pneumonia, bacteremia, meningitis, sinusitis, pleural empyema and endocarditis.

The polypeptide may be used as an antigen for vaccination of a host to produce specific antibodies which protect against invasion of bacteria, for example by blocking adherence of bacteria to damaged tissue. Examples of tissue damage include wounds in skin or connective tissue caused e.g. by mechanical, chemical or thermal damage or by implantation of indwelling devices, or wounds in the mucous membranes, such as the mouth, mammary glands, urethra or vagina.

The polypeptides or cells expressing them can be used as an immunogen to produce antibodies thereto. These antibodies can be, for example, polyclonal or monoclonal antibodies. The term antibodies also includes chimeric, single chain, and humanized antibodies, as well as Fab fragments, or the product of an Fab expression library. Various procedures known in the art may be used for the production of such antibodies and fragments.

Antibodies generated against the polypeptides of the present invention can be obtained by direct injection of the polypeptides into an animal or by administering the polypeptides to an animal, preferably a nonhuman. The antibody so obtained will then bind the polypeptides itself. In this manner, even a sequence encoding only a fragment of the polypeptides can be used to generate antibodies binding the whole native polypeptides. Such antibodies can then be used to isolate the polypeptide from tissue expressing that polypeptide.

Polypeptide derivatives include antigenically or immunologically equivalent derivatives which form a particular aspect of this invention.

The term 'antigenically equivalent derivative' as used herein encompasses a polypeptide or its equivalent which will be specifically recognised by certain antibodies which, when raised to the protein or polypeptide according to the present invention, interfere with the immediate physical interaction between pathogen and mammalian host.

The term 'immunologically equivalent derivative' as used herein encompasses a peptide or its equivalent which when used in a suitable formulation to raise antibodies in a vertebrate, the antibodies act to interfere with the immediate physical interaction between pathogen and mammalian host.

The polypeptide, such as an antigenically or immunologically equivalent derivative or a fusion protein thereof is used as an antigen to immunize a mouse or other animal such as a rat or chicken. The fusion protein may provide stability to the polypeptide. The antigen may be associated, for example by conjugation, with an immunogenic carrier protein for example bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). Alternatively a multiple antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically or immunologically equivalent polypeptide thereof may be sufficiently antigenic to improve immunogenicity so as to obviate the use of a carrier.

For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler and Milstein, 1975, *Nature*, 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor et al., 1983, *Immunology Today* 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole, et al., 1985, in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96).

Techniques described for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptide products of this invention.

Using the procedure of Kohler and Milstein (1975 *Nature* 256, 495-497), antibody-containing cells from the immunised mammal are fused with myeloma cells to create hybridoma cells secreting monoclonal antibodies.

The hybridomas are screened to select a cell line with high binding affinity and favorable cross reaction with other streptococcal species using one or more of the original polypeptide and/or the fusion protein. The selected cell line is cultured to obtain the desired Mab.

Hybridoma cell lines secreting the monoclonal antibody are another aspect of this invention.

Alternatively phage display technology could be utilised to select antibody genes with binding activities towards the polypeptide either from repertoires of PCR amplified v-genes of lymphocytes from humans screened for possessing anti-Fbp or from naive libraries (McCafferty, J. et al., (1990), *Nature* 348, 552-554; Marks, J. et al., (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. et al., (1991) *Nature* 352, 624-628).

The antibody should be screened again for high affinity to the polypeptide and/or fusion protein.

As mentioned above, a fragment of the final antibody may be prepared.

The antibody may be either intact antibody of  $M_r$  approx 150,000 or a derivative of it, for example a Fab fragment or a Fv fragment as described in Skerra, A and Pluckthun, A (1988) *Science* 240 1038-1040. If two antigen binding domains are present each domain may be directed against a different epitope - termed 'bispecific' antibodies.

The antibody of the invention may be prepared by conventional means for example by established monoclonal antibody technology (Kohler, G. and Milstein, C. (1975), *Nature*, 256, 495-497) or using recombinant means e.g. combinatorial libraries, for example as described in Huse, W.D. et al., (1989) *Science* 246, 1275-1281.

Preferably the antibody is prepared by expression of a DNA polymer encoding said antibody in an appropriate expression system such as described above for the expression of polypeptides of the invention. The choice of vector for the expression system will be determined in part by the host, which may be a prokaryotic cell, such as *E. coli* (preferably strain B) or *Streptomyces sp.* or a eukaryotic cell, such as a mouse C127, mouse myeloma, human HeLa, Chinese hamster ovary, filamentous or unicellular fungi or insect cell. The host may also be a transgenic animal or a transgenic plant (for example as described in Hiatt, A et al., (1989) *Nature* 34, 76-78). Suitable vectors include plasmids, bacteriophages, cosmids and recombinant viruses, derived from, for example, baculoviruses and vaccinia.

The Fab fragment may also be prepared from its parent monoclonal antibody by enzyme treatment, for example using papain to cleave the Fab portion from the Fc portion.

Preferably the antibody or derivative thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be 'humanised'; where the complementarity determining region(s) of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones, P. et al. (1986), *Nature* 321, 522-525 or Tempest et al., (1991) *Biotechnology* 9, 266-273.

The modification need not be restricted to one of 'humanisation'; other primate sequences (for example Newman, R. et al., 1992, *Biotechnology*, 10, 1455-1460) may also be used.

The humanised monoclonal antibody, or its fragment having binding activity, form a particular aspect of this invention.

This invention provides a method of screening drugs to identify those which interfere with the interaction of the cell surface protein or active fragment to mammalian cells, the method comprising incubating a mammalian cell or membrane preparation with labeled polypeptide in the presence of the drug and measuring the ability of the drug to block this interaction.

The use of a polynucleotide of the invention in DNA immunisation will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff et al., *Hum Mol Genet* 1992, 1:363, Manthorpe et al., *Hum. Gene Ther.* 1993:4, 419), delivery of DNA complexed with specific protein carriers (Wu et al., *J Biol Chem* 1989: 264,16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS*,1986:83,9551), encapsulation of DNA in various forms of liposomes (Kaneda et al., *Science* 1989:243,375), particle bombardment (Tang et al., *Nature* 1992, 356:152, Eisenbraun et al., *DNA Cell Biol* 1993, 12:791) and *in vivo* infection using cloned retroviral vectors (Seeger et al., *PNAS* 1984:81,5849). Suitable promoters for muscle transfection include CMV, RSV, SRA, actin, MCK, alpha globin, adenovirus and dihydrofolate reductase.

#### **IgAFcBP binding molecules and assays**

This invention also provides a method for identification of molecules, such as binding molecules, that bind IgAFcBP. Genes encoding proteins that bind IgAFcBP, such as binding proteins, can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Such methods are described in many laboratory manuals such as, for instance, Coligan et al., *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

For instance, expression cloning may be employed for this purpose. To this end polyadenylated RNA is prepared from a cell responsive to IgAFcBP, a cDNA library is created from this RNA, the library is divided into pools and the pools are transfected individually into cells that are not responsive to IgAFcBP. The transfected cells then are exposed to labeled IgAFcBP. IgAFcBP can be labeled by a variety of well-known techniques including standard methods of radio-iodination or inclusion of a recognition site for a site-specific protein kinase.) Following exposure, the cells are fixed and binding of IgAFcBP is determined. These procedures conveniently are carried out on glass slides.

Pools are identified of cDNA that produced IgAFcBP-binding cells. Sub-pools are prepared from these positives, transfected into host cells and screened as described above. Using an iterative sub-pooling and re-screening process, one or more single clones that encode the putative binding molecule, such as a binding molecule, can be isolated.

Alternatively a labeled ligand can be photoaffinity linked to a cell extract, such as a membrane or a membrane extract, prepared from cells that express a molecule that it binds, such as a binding molecule. Cross-linked material is resolved by polyacrylamide gel electrophoresis ("PAGE") and exposed to X-ray film. The labeled complex containing the ligand-binding can be excised, resolved into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing can be used to design unique or degenerate oligonucleotide probes to screen cDNA libraries to identify genes encoding the putative binding molecule.

Polypeptides of the invention also can be used to assess IgAFcBP binding capacity of IgAFcBP binding molecules, such as binding molecules, in cells or in cell-free preparations.

Polypeptides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics.

#### **Antagonists and Agonists - assays and molecules**

The invention also provides a method of screening compounds to identify those which enhance or block the action of IgAFcBP polypeptides or polynucleotides on cells, such as its interaction with binding molecules, such as IgAFcBP-binding molecules. An antagonist is a compound which decreases the natural biological functions of IgAFcBP. An agonist is a compound which increases the natural biological functions of IgAFcBP.

For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, may be prepared from a cell that expresses a molecule that binds IgAFcBP. The preparation is incubated with labeled IgAFcBP in the absence or the presence of a candidate molecule which may be a IgAFcBP agonist or antagonist. The ability of the candidate molecule to bind the binding molecule is reflected in decreased binding of the labeled ligand. Molecules which bind gratuitously, *i.e.*, without inducing the effects of IgAFcBP on binding the IgAFcBP binding molecule, are most likely to be good antagonists. Molecules that bind well and elicit effects that are the same as or closely related to IgAFcBP are agonists.

IgAFcBP-like effects of potential agonists and antagonists may be measured, for instance, by determining activity



of a reporter system following interaction of the candidate molecule with a cell or appropriate cell preparation, and comparing the effect with that of IgAFcBP or molecules that elicit the same effects as IgAFcBP. Reporter systems that may be useful in this regard include but are not limited to colorimetric labeled substrate converted into product, a reporter gene that is responsive to changes in IgAFcBP activity, and binding assays known in the art.

Another example of an assay for IgAFcBP antagonists is a competitive assay that combines IgAFcBP and a potential antagonist with membrane-bound IgAFcBP binding molecules, recombinant IgAFcBP binding molecules, natural substrates or ligands, or substrate or ligand mimetics, under appropriate conditions for a competitive inhibition assay. IgAFcBP can be labeled, such as by radioactivity or a colorimetric compound, such that the number of IgAFcBP molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to a polypeptide of the invention and thereby inhibit or extinguish its activity. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing IgAFcBP-induced activities, thereby preventing the action of IgAFcBP by excluding IgAFcBP from binding.

Potential antagonists include a small molecule which binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such as binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules.

Other potential antagonists include antisense molecules. Antisense technology can be used to control gene expression through antisense DNA or RNA or through double- or triple-helix formation. Antisense techniques are discussed, for example, in - Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance Lee et al., *Nucleic Acids Research* 6: 3073 (1979); Cooney et al., *Science* 241: 456 (1988); and Dervan et al., *Science* 251: 1360 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA. For example, the 5' coding portion of a polynucleotide that encodes the mature polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of IgAFcBP. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into IgAFcBP polypeptide. The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed in vivo to inhibit production of IgAFcBP.

In a particular aspect the invention provides the use of the polypeptide, polynucleotide or inhibitor of the invention to interfere with the immediate physical interaction between a pathogen and mammalian host responsible for sequelae of infection. In particular the molecules of the invention may be used: i) in the prevention of adhesion of bacteria, in particular gram positive bacteria, to mammalian extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; ii) to block cell surface protein mediated mammalian cell invasion by, for example, initiating phosphorylation of mammalian tyrosine kinases (Rosenshine et al., *Infect. Immun.* 60:2211 (1992)) iii) to block bacterial adhesion between mammalian extracellular matrix proteins and bacterial cell surface proteins which mediate tissue damage; iv) to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

Each of the DNA sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein upon expression can be used as a target for the screening of antibacterial drugs. Additionally, the DNA sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The antagonists and agonists of the invention may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as hereinafter described.

The antagonists and agonists may be employed for instance to inhibit conjunctivitis, sinusitis, adherence to and initiation of infection in the respiratory tract.

## Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal which comprises inoculating the individual with IgAFcBP, or a fragment or variant thereof, adequate to produce antibody to protect said individual from infection, particularly bacterial infection and most particularly streptococcal infection, such as *Streptococcus pneumoniae* infections. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises, through gene therapy, delivering gene encoding IgAFcBP, or a fragment or a variant thereof, for expressing IgAFcBP, or a fragment or a variant thereof in



*vivo* in order to induce an immunological response to produce antibody to protect said individual from disease.

A further aspect of the invention relates to an immunological composition which, when introduced into a host capable or having induced within it an immunological response, induces an immunological response in such host to a IgAFcBP or protein coded therefrom, wherein the composition comprises a recombinant IgAFcBP or protein coded therefrom comprising DNA which codes for and expresses an antigen of said IgAFcBP or protein coded therefrom.

The IgAFcBP or a fragment thereof may be fused with co-protein which may not by itself produce antibodies, but is capable of stabilizing the first protein and producing a fused protein which will have immunogenic and protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as Glutathione-S-transferase (GST) or beta-galactosidase, relatively large co-proteins which solubilise the protein and facilitate production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system. The co-protein may be attached to either the amino or carboxy terminus of the first protein.

Provided by this invention are methods using the described polynucleotide or particular fragments thereof that have been shown to encode non-variable regions of bacterial cell surface proteins. Also provided are DNA constructs comprising such fragments that are useful in such genetic immunization experiments in animal models of infection with *S. pneumoniae*. Such experiments will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value from the requisite organ of the animal successfully resisting or clearing infection for the development of prophylactic agents or therapeutic treatments of *S. pneumoniae* infection in mammals, particularly humans.

The present invention also includes a vaccine formulation which comprises the immunogenic recombinant protein together with a suitable carrier. Since the protein may be broken down in the stomach, it is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Whilst the invention has been described with reference to certain IgAFcBP, it is to be understood that this covers fragments of the naturally occurring protein and similar proteins with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant protein.

### Compositions

The invention also relates to compositions comprising the polynucleotide or the polypeptides discussed above or the agonists or antagonists. Thus, the polypeptides of the present invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to a subject. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration.

### Kits

The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, reflecting approval by the agency of the manufacture, use or sale of the product for human administration.

### Administration

Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance,

administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

The pharmaceutical compositions generally are administered in an amount effective for treatment or prophylaxis of a specific indication or indications. In general, the compositions are administered in an amount of at least about 10  $\mu\text{g/kg}$  body weight. In most cases they will be administered in an amount not in excess of about 8  $\text{mg/kg}$  body weight per day. Preferably, in most cases, dose is from about 10  $\mu\text{g/kg}$  to about 1  $\text{mg/kg}$  body weight, daily. It will be appreciated that optimum dosage will be determined by standard methods for each treatment modality and indication, taking into account the indication, its severity, route of administration, complicating conditions and the like.

In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

Alternatively the composition may be formulated for topical application for example in the form of ointments, creams, lotions, eye ointments, eye drops, ear drops, mouthwash, impregnated dressings and sutures and aerosols, and may contain appropriate conventional additives, including, for example, preservatives, solvents to assist drug penetration, and emollients in ointments and creams. Such topical formulations may also contain compatible conventional carriers, for example cream or ointment bases, and ethanol or oleyl alcohol for lotions. Such carriers may constitute from about 1% to about 98% by weight of the formulation; more usually they will constitute up to about 80% by weight of the formulation.

For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01  $\text{mg/kg}$  to 10  $\text{mg/kg}$ , typically around 1  $\text{mg/kg}$ . The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

In-dwelling devices include surgical implants, prosthetic devices and catheters, i.e., devices that are introduced to the body of an individual and remain in position for an extended time. Such devices include, for example, artificial joints, heart valves, pacemakers, vascular grafts, vascular catheters, cerebrospinal fluid shunts, urinary catheters, continuous ambulatory peritoneal dialysis (CAPD) catheters, etc.

The composition of the invention may be administered by injection to achieve a systemic effect against relevant bacteria shortly before insertion of an in-dwelling device. Treatment may be continued after surgery during the in-body time of the device. In addition, the composition could also be used to broaden perioperative cover for any surgical technique to prevent streptococcal wound infections.

Many orthopaedic surgeons consider that humans with prosthetic joints should be considered for antibacterial prophylaxis before dental treatment that could produce a bacteraemia. Late deep infection is a serious complication sometimes leading to loss of the prosthetic joint and is accompanied by significant morbidity and mortality. It may therefore be possible to extend the use of the active agent as a replacement for prophylactic antibacterials in this situation.

In addition to the therapy described above, the compositions of this invention may be used generally as a wound treatment agent to prevent adhesion of bacteria to matrix proteins exposed in wound tissue and for prophylactic use in dental treatment as an alternative to, or in conjunction with, antibacterial prophylaxis.

Alternatively, the composition of the invention may be used to bathe an indwelling device immediately before insertion. The active agent will preferably be present at a concentration of 1  $\mu\text{g/ml}$  to 10  $\text{mg/ml}$  for bathing of wounds or indwelling devices.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response.

A suitable unit dose for vaccination is 0.5-5  $\mu\text{g/kg}$  of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks.

With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

The antibodies described above may also be used as diagnostic reagents to detect the presence of bacteria containing the cell surface protein.

In order to facilitate understanding of the following example certain frequently occurring methods and/or terms will be described.

### Gene therapy

The IgAFcBP polynucleotides, polypeptides, agonists and antagonists that are polypeptides may be employed in accordance with the present invention by expression of such polypeptides *in vivo*, in treatment modalities often referred to as "gene therapy". The compounds of the inventions may be used as gene immunotherapies, to engender an immune response in an individual against the organism from which such compound was derived as well as related organisms.

Thus, for example, cells from a individual may be engineered with a polynucleotide, such as a DNA or RNA, encoding a polypeptide *ex vivo*, and the engineered cells then can be provided to an individual to be treated with the polypeptide. For example, cells may be engineered *ex vivo* by the use of a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention. Such methods are well-known in the art and their use in the present invention will be apparent from the teachings herein.

Similarly, cells may be engineered *in vivo* for expression of a polypeptide *in vivo* by procedures known in the art. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct then may be isolated and introduced into a packaging cell is transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to an individual for engineering cells *in vivo* and expression of the polypeptide *in vivo*. These and other methods for administering a polypeptide of the present invention by such method should be apparent to those skilled in the art from the teachings of the present invention.

Retroviruses from which the retroviral plasmid vectors herein above mentioned may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, adenovirus, Myeloblastic Sarcoma Virus, and mammary tumor virus. In one embodiment, the retroviral plasmid vector is derived from Moloney Murine Leukemia Virus.

Such vectors well include one or more promoters for expressing the polypeptide. Suitable promoters which may be employed include, but are not limited to, the retroviral LTR; the SV40 promoter; and the human cytomegalovirus (CMV) promoter described in Miller et al., *Biotechniques* 7: 980-990 (1989), or any other promoter (e.g., cellular promoters such as eukaryotic cellular promoters including, but not limited to, the histone, RNA polymerase III, and  $\beta$ -actin promoters). Other viral promoters which may be employed include, but are not limited to, adenovirus promoters, thymidine kinase (TK) promoters, and B19 parvovirus promoters. The selection of a suitable promoter will be apparent to those skilled in the art from the teachings contained herein.

The nucleic acid sequence encoding the polypeptide of the present invention will be placed under the control of a suitable promoter. Suitable promoters which may be employed include, but are not limited to, adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the RSV promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoA1 promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs (including the modified retroviral LTRs herein above described); the  $\beta$ -actin promoter; and human growth hormone promoters. The promoter also may be the native promoter which controls the gene encoding the polypeptide.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, Y-2, Y-AM, PA12, T19-14X, VT-19-17-H2, YCRE, YCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, A., *Human Gene Therapy* 1: 5-14 (1990). The vector may be transduced into the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and  $\text{CaPO}_4$  precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

The producer cell line will generate infectious retroviral vector particles, which include the nucleic acid sequence (s) encoding the polypeptides. Such retroviral vector particles then may be employed to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express the nucleic acid sequence(s) encoding the polypeptide. Eukaryotic cells which may be transduced include, but are not limited to, embryonic stem cells, embryonic carcinoma cells, as well as hematopoietic stem cells, hepatocytes, fibroblasts, myoblasts, keratinocytes, endothelial cells, and bronchial epithelial cells.

## EXAMPLES

The present invention is further described by the following examples. The examples are provided solely to illustrate the invention by reference to specific embodiments. These exemplification's, while illustrating certain specific aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

Certain terms used herein are explained in the foregoing glossary.

All examples were carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. Routine molecular biology techniques of the following examples can be carried out as described in standard laboratory manuals, such as Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

All parts or amounts set out in the following examples are by weight, unless otherwise specified.

Unless otherwise stated size separation of fragments in the examples below was carried out using standard techniques of agarose and polyacrylamide gel electrophoresis ("PAGE") in Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) and numerous other references such as, for instance, by Goeddel et al., *Nucleic Acids Res.* 8: 4057 (1980).

Unless described otherwise, ligations were accomplished using standard buffers, incubation temperatures and times, approximately equimolar amounts of the DNA fragments to be ligated and approximately 10 units of T4 DNA ligase ("ligase") per 0.5 µg of DNA.

The polynucleotide having the DNA sequence given in SEQ ID NO: 1 was obtained from the sequencing of a library of clones of chromosomal DNA of *Streptococcus pneumoniae* 0100993 in *E. coli*. It has been demonstrated by the process herein described that it is transcribed *in vivo* in an established infection of *Streptococcus pneumoniae* 0100993 in a mouse model of infection.

To obtain the polynucleotide encoding the cell surface protein using the DNA sequence given in SEQ ID NO: 1 typically a library of clones of chromosomal DNA of *Streptococcus pneumoniae* 0100993 in *E. coli* or some other suitable host is probed with a radiolabelled oligonucleotide, preferably a 17mer or longer, derived from the partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using high stringency washes. By sequencing the individual clones thus identified with sequencing primers designed from the original sequence it is then possible to extend the sequence in both directions to determine the full gene sequence. Conveniently such sequencing is performed using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see Screening By Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70).

#### Example 1 Isolation of DNA coding for Novel IgAFcBP from *S. pneumoniae*

The polynucleotide having the DNA sequence given in SEQ ID NO:1 was obtained from a library of clones of chromosomal DNA of *Streptococcus pneumoniae* in *E. coli*. In some cases the sequencing data from two or more clones containing overlapping *Streptococcus pneumoniae* DNA was used to construct the contiguous DNA sequence in SEQ ID NO:1. Libraries may be prepared by routine methods, for example: Methods 1 and 2 below.

Total cellular DNA is isolated from *Streptococcus pneumoniae* strain 0100993 according to standard procedures and size-fractionated by either of two methods.

##### Method 1

Total cellular DNA is mechanically sheared by passage through a needle in order to size-fractionate according to standard procedures. DNA fragments of up to 11kbp in size are rendered blunt by treatment with exonuclease and DNA polymerase, and EcoRI linkers added. Fragments are ligated into the vector Lambda ZapII that has been cut with EcoRI, the library packaged by standard procedures and *E. coli* infected with the packaged library. The library is amplified by standard procedures.

##### Method 2

Total cellular DNA is partially hydrolyzed with a combination of four restriction enzymes (RsaI, PstI, AluI and Bsh1235I) and size-fractionated according to standard procedures. EcoRI linkers are ligated to the DNA and the fragments then ligated into the vector Lambda ZapII that have been cut with EcoRI, the library packaged by standard procedures, and *E. coli* infected with the packaged library. The library is amplified by standard procedures.

#### Example 2 Gene immunotherapeutic expression of *S. pneumoniae* IgAFcBP

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature overnight. After 24 hours at room temperature, the flask is inverted - the chunks of tissue remain fixed to the bottom of the flask - and fresh media is added (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin). The tissue is then incubated at 37°C for approximately one week. At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerges. The monolayer is trypsinized and scaled into larger flasks.

A vector for gene therapy is digested with restriction enzymes for cloning a fragment to be expressed. The digested

vector is treated with calf intestinal phosphatase to prevent self-ligation. The dephosphorylated, linear vector is fractionated on an agarose gel and purified.

IgAFcBP DNA or a fragment thereof capable of expressing active IgAFcBP or a fragment thereof, is isolated. The ends of the fragment are modified, if necessary, for cloning into the vector. For instance, 5' overhanging may be treated with DNA polymerase to create blunt ends. 3' overhanging ends may be removed using S1 nuclease. Linkers may be ligated to blunt ends with T4 DNA ligase.

Equal quantities of the Moloney murine leukemia virus linear backbone and the IgAFcBP fragment are mixed together and joined using T4 DNA ligase. The ligation mixture is used to transform *E. coli* and the bacteria are then plated onto agar-containing kanamycin. Kanamycin phenotype and restriction analysis confirm that the vector has the properly inserted gene.

Packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The vector containing the IgAFcBP gene is introduced into the packaging cells by standard techniques. Infectious viral particles containing the IgAFcBP gene are collected from the packaging cells, which now are called producer cells.

Fresh media is added to the producer cells, and after an appropriate incubation period media is harvested from the plates of confluent producer cells. The media, containing the infectious viral particles, is filtered through a Millipore filter to remove detached producer cells. The filtered media then is used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the filtered media. Polybrene (Aldrich) may be included in the media to facilitate transduction. After appropriate incubation, the media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his, to select out transduced cells for expansion.

Engineered fibroblasts then may be injected into mice or rats, either alone or after having been grown to confluence on microcarrier beads, such as cytodex 3 beads. The injected fibroblasts produce IgAFcBP product, and the biological actions of the protein, most particularly the induction of an immune response, are conveyed to the host.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

SEQUENCE LISTING

5

(1) GENERAL INFORMATION

10

(i) APPLICANT: SmithKline Beecham Corporation and  
SmithKline Beecham p.l.c.

15

(ii) TITLE OF THE INVENTION: Novel IgA Fc Binding Protein

20

(iii) NUMBER OF SEQUENCES: 4

25

(iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: SmithKline Beecham, Corporate  
Intellectual Property  
(B) STREET: Two New Horizons Court  
(C) CITY: Brentford  
(D) STATE: Middlesex  
(E) COUNTRY: United Kingdom  
(F) POST CODE: TW8 9EP

30

(v) COMPUTER READABLE FORM:

35

(A) MEDIUM TYPE: Diskette  
(B) COMPUTER: IBM Compatible  
(C) OPERATING SYSTEM: DOS  
(D) SOFTWARE: FastSEQ for Windows Version 2.0

40

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:

45

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 60/027,030  
(B) FILING DATE: 24-SEP-1996

50

(A) APPLICATION NUMBER: 60/040,656  
(B) FILING DATE: 10-MAR-1997

55

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: CONNELL, Anthony Christopher  
(B) REGISTRATION NUMBER: 5630 and 26758

(C) REFERENCE/DOCKET NUMBER: P50545

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: +44 127 964 4395

(B) TELEFAX: +44 181 975 6294

(C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1056 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTGGGAGACA TTAAGCGAAC GTATTTGTAT AAATTAGATG AATCAACCCA  
AAAAGCCCAA 60  
CTACAGGAAC TGGTCACAGA AAGTCAATCA AAAGTAGATG AAGCTTTTTTC  
TAAATTTAAA 120  
AATGGCTTAT CTTCTTCGTC AAGTTCAGGA TCCTCCACTA AACCAGAAAC  
TCCGCAGCCG 180  
GAAAAACCAG AGCATCAAAA ACCAACAACCT CCAGCTCCGG ATACCAAACC  
AAGCCCTCAA 240  
CCAGAAGGCA AGAAACCAAG CGTACCAGAC ATTAATCAGG AAAAAGAAAA  
AGCTAAGCTT 300  
GCTGTAGCAA CCTACATGAG CAAGATTTTA GATGATATAC AAAACATCA  
TCTGCAGAAA 360  
GAAAAACATC GTCAGATTGT TGCTCTTATT AAGGAGCTTG ATGAGCTTAA  
AAAGCAAGCT 420  
CTTTCTGAAA TTGATAATGT AAATACCAAA GTAGAAATTG AAAATACAGT  
CCACAAGATA 480  
TTTGCAGACA TGGATGCAGT TGTGACTAAA TTCAAAAAAG GCTTAACTCA  
GGACACACCA 540  
AAAGAACCAG GTAACAAAAA ACCATTTGCT CCAAAACCAG GTATGCAACC  
AAGTCCTCAA 600  
CCAGAGGTTA AACCGCAGCT AGAAAAACCA AAACCAGAGG TTAACCSCA  
ACCAGAAAAA 660  
CCAAAAACCAG AGGTTAAACC GCAACCAGAA AAACCAAAAC CAGAGGTTAA  
ACCGCAGCCG 720  
GAAAAACCAA AACCAGAGST TAAACCGCAA CCAGAAAAAC CAAAACCAGA  
GGTTAAACCG 780



CAGCCGAAAA AACCAAAACC AGAGGTTAAA CCGCAGCTGG AAAAACCAAA  
 ACCAGATAAT 840  
 5 AGCAAGCCAC AAGCAGATGA TAAGAAGCCA TCAACTACAA ATAATTTAAG  
 CAAGGACAAG 900  
 CAACCTTCTA ACCAAGCTTC AACAAACGAA AAAGCAACAA ATAAACCGAA  
 GAAGTCATTG 960  
 10 CCATCAACTG GATCTATTTC AAATCTAGCA CTTGAAATTG CAGGTCTTCT  
 TACCTTGGCG 1020  
 GGGGCAACCA TTCTTGCTAA GAAAAGAATG AAATAG  
 1056

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 351 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Leu Gly Asp Ile Lys Arg Thr Tyr Leu Tyr Lys Leu Asp Glu Ser Thr  
 1 5 10 15  
 30 Gln Lys Ala Gln Leu Gln Glu Leu Val Thr Glu Ser Gln Ser Lys Leu  
 20 25 30  
 Asp Glu Ala Phe Ser Lys Phe Lys Asn Gly Leu Ser Ser Ser Ser  
 35 40 45  
 Ser Gly Ser Ser Thr Lys Pro Glu Thr Pro Gln Pro Glu Lys Pro Glu  
 50 55 60  
 His Gln Lys Pro Thr Thr Pro Ala Pro Asp Thr Lys Pro Ser Pro Gln  
 65 70 75 80  
 40 Pro Glu Gly Lys Lys Pro Ser Val Pro Asp Ile Asn Gln Glu Lys Glu  
 85 90 95  
 Lys Ala Lys Leu Ala Val Ala Thr Tyr Met Ser Lys Ile Leu Asp Asp  
 100 105 110  
 45 Ile Gln Lys His His Leu Gln Lys Glu Lys His Arg Gln Ile Val Ala  
 115 120 125  
 Leu Ile Lys Glu Leu Asp Glu Leu Lys Lys Gln Ala Leu Ser Glu Ile  
 130 135 140  
 Asp Asn Val Asn Thr Lys Val Glu Ile Glu Asn Thr Val His Lys Ile  
 50 145 150 155 160  
 Phe Ala Asp Met Asp Ala Val Val Thr Lys Phe Lys Lys Gly Leu Thr  
 165 170 175  
 Gln Asp Thr Pro Lys Glu Pro Gly Asn Lys Lys Pro Phe Ala Pro Lys  
 55 180 185 190

Pro Gly Met Gln Pro Ser Pro Gln Pro Glu Val Lys Pro Gln Leu Glu  
 195 200 205  
 5 Lys Pro Lys Pro Glu Val Lys Pro Gln Pro Glu Lys Pro Lys Pro Glu  
 210 215 220  
 Val Lys Pro Gln Pro Glu Lys Pro Lys Pro Glu Val Lys Pro Gln Pro  
 225 230 235 240  
 10 Glu Lys Pro Lys Pro Glu Val Lys Pro Gln Pro Glu Lys Pro Lys Pro  
 245 250 255  
 Glu Val Lys Pro Gln Pro Glu Lys Pro Lys Pro Glu Val Lys Pro Gln  
 260 265 270  
 15 Leu Glu Lys Pro Lys Pro Asp Asn Ser Lys Pro Gln Ala Asp Asp Lys  
 275 280 285  
 Lys Pro Ser Thr Thr Asn Asn Leu Ser Lys Asp Lys Gln Pro Ser Asn  
 290 295 300  
 20 Gln Ala Ser Thr Asn Glu Lys Ala Thr Asn Lys Pro Lys Lys Ser Leu  
 305 310 315 320  
 Pro Ser Thr Gly Ser Ile Ser Asn Leu Ala Leu Glu Ile Ala Gly Leu  
 325 330 335  
 25 Leu Thr Leu Ala Gly Ala Thr Ile Leu Ala Lys Lys Arg Met Lys  
 340 345 350

## (2) INFORMATION FOR SEQ ID NO:3:

30 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 35 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

40 ATTAAGCGAA CGTATTTGTA  
 20

## (2) INFORMATION FOR SEQ ID NO:4:

45 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 base pairs  
 (B) TYPE: nucleic acid  
 50 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

55

GGAATTTAGT CACAAC TGCA

20

5

**Claims**

10

1. An isolated polynucleotide comprising a member selected from the group consisting of:

(a) a polynucleotide having at least a 70% identity to a polynucleotide encoding a polypeptide comprising amino acids 1 to 170 of SEQ ID NO:2;  
 (b) a polynucleotide which is complementary to the polynucleotide of (a); and  
 (c) a polynucleotide comprising at least 15 bases of the polynucleotide of (a) or (b).

15

2. The polynucleotide of Claim 1 wherein the polynucleotide is DNA.

3. The polynucleotide of Claim 1 wherein the polynucleotide is RNA.

20

4. The polynucleotide of Claim 2 comprising nucleotide 1 to 510 set forth in SEQ ID NO:1.

5. The polynucleotide of Claim 2 comprising a sequence encoding the amino acid sequence of SEQ ID NO:2 as set forth in SEQ ID NO:1.

25

6. The polynucleotide of Claim 2 which encodes a polypeptide comprising amino acid 1 to 170 of SEQ ID NO:2.

7. An isolated polynucleotide comprising a member selected from the group consisting of:

(a) a polynucleotide having at least a 70% identity to a polynucleotide encoding the same mature polypeptide expressed by DNA isolated from NCIMB Deposit No. 40794;  
 (b) a polynucleotide complementary to the polynucleotide of (a); and  
 (c) a polynucleotide comprising at least 15 bases of the polynucleotide of (a) or (b).

30

8. A vector comprising the DNA of Claim 2.

35

9. A host cell comprising the vector of Claim 8.

10. A process for producing a polypeptide comprising: expressing from the host cell of Claim 9 a polypeptide encoded by said DNA.

40

11. A process for producing a cell which expresses a polypeptide comprising transforming or transfecting the cell with the vector of Claim 8 such that the cell expresses the polypeptide encoded by the cDNA contained in the vector.

12. A polypeptide comprising an amino acid sequence which is at least 70% identical to amino acid 1 to 170 of SEQ ID NO:2.

45

13. A polypeptide comprising an amino acid sequence as set forth in SEQ ID NO:2.

14. An antibody against the polypeptide of Claim 12.

50

15. An antagonist which inhibits the activity of the polypeptide of Claim 12.

16. A method for the treatment of an individual having need of IgAFcBP comprising: administering to the individual a therapeutically effective amount of the polypeptide of Claim 12.

55

17. The method of Claim 16 wherein said therapeutically effective amount of the polypeptide is administered by providing to the individual DNA encoding said polypeptide and expressing said polypeptide *in vivo*.

18. A method for the treatment of an individual having need to inhibit IgAFcBP polypeptide comprising: administering to the individual a therapeutically effective amount of the antagonist of Claim 15.

19. A process for diagnosing a disease related to expression of the polypeptide of Claim 12 comprising:  
determining a nucleic acid sequence encoding said polypeptide.

20. A diagnostic process comprising:  
analyzing for the presence of the polypeptide of Claim 12 in a sample derived from a host.

21. A method for identifying compounds which bind to and inhibit an activity of the polypeptide of Claim 12 comprising:  
contacting a cell expressing on the surface thereof a binding for the polypeptide, said binding being associated with a second component capable of providing a detectable signal in response to the binding of a compound to said binding, with a compound to be screened under conditions to permit binding to the binding; and  
determining whether the compound binds to and activates or inhibits the binding by detecting the presence or absence of a signal generated from the interaction of the compound with the binding.

22. A method for inducing an immunological response in a mammal which comprises inoculating the mammal with IgAFcBP, or a fragment or variant thereof, adequate to produce antibody to protect said animal from disease.

23. A method of inducing immunological response in a mammal which comprises, through gene therapy, delivering gene encoding IgAFcBP fragment or a variant thereof, for expressing IgAFcBP, or a fragment or a variant thereof *in vivo* in order to induce an immunological response to produce antibody to protect said animal from disease.

24. An immunological composition which, when introduced into a mammalian host, induces an immunological response in that mammal to a given IgAFcBP polynucleotide or protein coded therefrom, wherein the composition comprises a recombinant IgAFcBP polynucleotide or protein coded therefrom comprising DNA which codes for and expresses an antigen of said IgAFcBP polynucleotide or protein coded therefrom.

FIGURE 1. IgAFcBP cDNA Sequence [SEQ ID NO:1]

```
1  ATTAAGCGAA CGTATTTGTA TAAATTAGAT GAATCAACCC AAAAAGCCCA
51  ACTACAGGAA CTGGTCACAG AAAGTCAATC AAAACTAGAT GAAGCTTTTN
101 CTAAATTTAA AAATGGCTTA TCTTCTTCGT CAAGTTCAGG ATCCTCCACT
151 AAACCAGAAA CTCCGCAGCC GGAAAAACCA GAGCATCAAA AACCAACAAC
201 TCCAGCTCCG GATACCAAAC CAAGCCCTCA ACCAGAAGGC AAGAAACCAA
251 GCGTACCAGA CATTAATCAG GAAAAAGAAA AAGCTAAGCT TGCTGTAGCA
301 ACCTACATGA GCAAGATTTT AGATGATATA CAAAACATC ATCTGCAGAA
351 AGAAAAACAT CGTCAGATTG TTGCTCTTAT TAAGGAGCTT GATGAGCTTA
401 AAAAGCAAGC TCTTTCTGAA ATTGATAATG TAAATACCAA AGTAGAAATT
451 GAAAATACAG TCCACAAGAT ATTTGCAGAC ATGGATGCAG TTGTGACTAA
501 ATTCCAAAAA
```

**FIGURE 2. IgAFcBP predicted amino acid sequence [SEQ ID NO:2]**

```
1  IKRTYLYKLD ESTQKAQLQE LVTESQSKLD EAFXKFKNGL SSSSSSGSST
51  KPETPQPEKP EHQKPTTPAP DTKPSPQPEG KKPSVPDINQ EKEKAKLAVA
101 TYMSKILDDI QKHHLQKEKH RQIVALIKEL DELKKQALSE IDNVNTKVEI
151 ENTVHKIFAD MDAVVTKFQK
```

**This Page Blank (uspto)**